



# Genetic Divergence Analysis in Amaranth (*Amaranthus spp.*) by Using Morpho-agronomic Traits in Western Uttar Pradesh, India

Harshit Tomar <sup>a\*</sup>, Vipin Kumar <sup>a</sup>, Rupesh Kumar <sup>a</sup>,  
Sudhanshu Singh <sup>a</sup>, Satya Prakash <sup>a</sup>, Yashraj <sup>a</sup>  
and Vikash Tyagi <sup>a</sup>

<sup>a</sup> Department of Vegetable Science, College of Horticulture, Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut-250 110 (U.P.), India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i92323

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100708>

Original Research Article

Received: 01/04/2023

Accepted: 01/06/2023

Published: 13/07/2023

## ABSTRACT

The field experiment was conducted during 2021-22. During the study the analysis of variance for 19 diverse genotypes of *Amaranthus* were designed in randomized block design and revealed significant difference for all the 12 characters, this indicated the presence of wide spectrum of variability among the genotypes. The analysis of genetic divergence through Mahalanobis  $D^2$  statistics revealed that a considerable genetic diversity was found among genotypes. 19 genotypes of *Amaranthus* were grouped into 5 clusters. Out of these 5 clusters, Cluster II contains maximum number of genotypes that is seven, followed by cluster I which comprising six genotypes, Cluster III comprising four genotypes, whereas cluster IV and V containing only one genotype each. All the

\*Corresponding author: E-mail: [tomarharshit180@gmail.com](mailto:tomarharshit180@gmail.com);

desired combinations of traits should be considered, while breeding programme for selecting high yielding genotypes and suitable for breeders to achieving improved plant type.

**Keywords:** Genetic diversity; *Amaranthus*; morpho-agronomic traits; breeding.

## 1. INTRODUCTION

"*Amaranthus* is a genus belonging to the Amaranthaceae family that has originated in South America. This genus contains approximately 70 species and out of these most of them are cultivated as leafy vegetables, grains and ornamental plants in different parts of the world" [1,2]. "There are three major grain producing *Amaranthus* species, *A. caudatus*, *A. cruentus* and *A. hypochondriacus*, all believed to originate from Central and South America; and three major leafy vegetable species, *A. tricolor*, *A. dubius* and *A. blitum* (*A. lividus*), of which *A. tricolor* is thought to originate from India or Southern China, *A. blitum* from Central Europe and *A. dubius* from Central America" [3]. "*Amaranthus* species are significant food crops that can withstand heat, drought, diseases, and pests" [4]. "It is one of those rare plants whose leaves are eaten as a vegetable while the seeds are used as cereals. Besides, it is also used as fodder, ornamental, organic red dye and for industrial purposes. It is high in protein, particularly in the amino acid, Lysine, which seeds is low in the cereal grains. Amaranth is one of the highest grains in fiber content. This makes Amaranth an effective agent against cancer and heart disease. Amaranth is also rich in many vitamins and minerals and it also does not contains gluten. They grow in a wide range of agroecological zones and are found in most tropical and subtropical areas" [5,6]. In India it is cultivated in Jammu and Kashmir, Himachal Pradesh, Uttarakhand, Sikkim, Assam, Meghalaya, Arunachal Pradesh, Nagaland, Tripura, Jharkhand, Chhattisgarh, Maharashtra, Gujarat, Orissa, Karnataka, Kerala and Tamil Nadu both in hill and plain regions.

The grain *Amaranthus* are paleo-allote tetraploids. The multivariate analysis has been established by several investigators for measuring the degree of divergence and for ascertaining the relative contribution of different characters to the total divergence. Wide genotypic variation has been seen in *Amaranthus* as a result of frequent interspecific and intervarietal hybridization. *Amaranthus* also exhibit a tremendous level of diversity. To assess

the genetic diversity of local *Amaranthus*, it is crucial to determine the proper genotype. Maintaining genetic variety, examining local genetic diversity, and selecting ecotypes with high nutritional value in their native environments all require the identification and preservation of germplasm. *Amaranthus* is still a minor crop that is not fully utilised for vegetable production. Despite having outstanding nutritional properties, little work has been done to improve its genetic profile.

*Amaranthus* shows a wide diversity in growth habit, leaf shape, colour and size, plant size and inflorescence characteristics. Among several statistical methods developed for measuring divergence between populations, multivariate analysis of D2 statistics has been effectively used for quantitative estimation of genetic variability according to Mahalanobis [7] D2 statistics, which can be effectively used for assessing the genetic divergence between populations and helping in selection of desirable parents for crossing programme. Recent research indicates that "under cultivated conditions, Amaranth produces fresh leaf yields of up to 40 t/ha. The yield of grain amaranth is highly variable with 1000 kg/ha considered a good yield" [4]. The purpose of this research is to Genetic Divergence Analysis in Amaranth (*Amaranthus* spp.) by using Morpho-agronomic Traits in Western Uttar Pradesh, India.

## 2. MATERIALS AND METHODS

The experiment was conducted during 2021-22 at the Horticultural Research Center of the Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, Uttar Pradesh, using 19 diverse germplasm collected from IARI New Delhi as shown in Table 1. Randomized block design with three replications was used to conduct the experiment. The observations were recorded on randomly selected plants for the various characters i.e. days to germination, days to 50% flowering, days to maturity, plant height (cm), inflorescence length (cm), number of leaves per plant, number of branches per plant, fresh leaf weight (g), biological yield per plant (g), biological yield ton per hectare, seed yield per plant (g) and seed yield per hectare (kg/ha).

**Table 1. List of cultivars/Germplasm included in the trial**

S. No.	Notation	Germplasm	Source
1	T <sub>1</sub>	Arun	Division of Vegetable Science, IARI, New Delhi
2	T <sub>2</sub>	Arka Samraksha	IIHR, Bengaluru
3	T <sub>3</sub>	Arka Suguma	IIHR, Bengaluru
4	T <sub>4</sub>	CO-2	Coimbatore, Tamilnadu
5	T <sub>5</sub>	CO-3	Coimbatore, Tamilnadu
6	T <sub>6</sub>	CO-5	Coimbatore, Tamilnadu
7	T <sub>7</sub>	Krishna Sree	Division of Vegetable Science, IARI, New Delhi
8	T <sub>8</sub>	IC-151606	Division of Vegetable Science, IARI, New Delhi
9	T <sub>9</sub>	IIHR-109-1	IIHR, Bengaluru
10	T <sub>10</sub>	RNA-1	Division of Vegetable Science, IARI, New Delhi
11	T <sub>11</sub>	Reni Sree	Division of Vegetable Science, IARI, New Delhi
12	T <sub>12</sub>	Pusa Lal Chaulai	Division of Vegetable Science, IARI, New Delhi
13	T <sub>13</sub>	Pusa Kiran	Division of Vegetable Science, IARI, New Delhi
14	T <sub>14</sub>	Arka Arunima	IIHR, Bengaluru
15	T <sub>15</sub>	Arka Verna	IIHR, Bengaluru
16	T <sub>16</sub>	CO-4	Coimbatore, Tamilnadu
17	T <sub>17</sub>	Kannara Local	Division of Vegetable Science, IARI, New Delhi
18	T <sub>18</sub>	IIHR-109-4	IIHR, Bengaluru
19	T <sub>19</sub>	IC-151608	Division of Vegetable Science, IARI, New Delhi

## 2.1 Statistical Analysis

All the statistical analysis to assess the genetic diversity in *Amaranthus* were performed by using standard formulas.

## 3. RESULTS AND DISCUSSION

Significant differences were found for all the twelve characters in the analysis of variance for the 19 genotypes, revealing a wide range of variability among the genotypes.

All 19 genotypic sample of *Amaranthus* under research were divided into 5 clusters based on Mahalanobis D<sup>2</sup> values. The distributions of *Amaranthus* genotypes into five clusters are presented in Table 2. Cluster II contains maximum number of 07 genotypes comprising namely as Arun, Arka Samraksha, CO-2, CO-3, Krishna Sree, Reni Sree and Kannara Local, followed by cluster I which comprising 06 genotypes namely Arka Suguma, CO-5, IIHR-109-1, RNA-1, Arka Arunima and Pusa Kiran, Cluster III comprising 04 genotypes namely IC-151606, Arka Verna, CO-4 and IIHR-109-4, whereas cluster IV and V containing only 01 genotype in each, namely IC-151608 and Pusa Lal Chaulai respectively. The small variation in D<sup>2</sup> values makes it clear that the genotypes in the cluster do not differ significantly in terms of their relative genetic distance. Therefore, these genotypes that are genetically different can be used as promising genotypes for use as parents in hybridization.

Based on the statistically significant difference in cluster means for several parameters, diversity among the genotypes was also assessed. The cluster mean calculated for twelve characters under study have been presented in Table 3. Days to germination showed highest mean for cluster number IV (7.07) followed by cluster V (7.00), whereas lowest mean in cluster number I (6.11). 50 percent flowering revealed highest mean in cluster number IV (57.47) followed by cluster III (52.70) and lowest mean for cluster number I (46.18), Days to maturity exhibited highest mean in cluster number IV (104.93) and lowest mean in cluster number II (96.08). Plant height revealed highest mean in cluster number IV (71.21) and lowest mean in cluster number II (51.27). Inflorescence length showed highest mean in cluster number III (33.16) and lowest mean in cluster number V (25.93). Number of leaves per plant exhibited highest mean in cluster number IV (45.07) and lowest mean in cluster number III (36.09). Number of branches per plant revealed highest mean in cluster number IV (3.80) and lowest in cluster V (2.20). Fresh leaf weight exhibited highest mean in cluster number IV (6.05) and lowest mean in cluster number V (3.36). Biological yield per plant revealed highest mean in cluster number V (212.82) and lowest mean in cluster number II (123.00). Biological yield ton per ha exhibited highest mean in cluster number V (45.14) and lowest mean in cluster number II (16.85). Seed yield kg per ha showed highest mean in cluster number V (15.91) and lowest mean in cluster

**Table 2. Number of genotypes in each cluster**

Clusters	No of genotypes	Genotypes
I	6	Arka Suguna, CO-5, IIHR-109-1, RNA-1, Arka Arunima, Pusa Kiran
II	7	Arun, Arka Samraksha, CO-2, CO-3, Krishna Sree, Reni Sree, Kannara local
III	4	IC-151606, Arka Verna, CO-4, IIHR-109-4
IV	1	IC-151608
V	1	Pusa Lal Chulai

**Table 3. Cluster mean of 19 genotypes of Amaranth for 12 characters**

Clusters		Days to Germination	50% Flowering	Days to maturity	Plant height (cm)	Inflorescence length (cm)	No. of leaf per plant	No. of branches per plant	Fresh leaf weight (g)	Biological yield per plant (g)	Biological yield ton/ha	Seed yield per ha (kg/ha.)	Seed yield per plant (g)
I	<b>Mean</b>	6.11	46.18	96.26	59.14	31.31	38.52	2.55	3.79	193.68	28.28	7.37	159.69
	<b>± SE</b>	0.43	3.96	4.92	6.94	1.19	5.18	0.28	0.53	6.66	3.12	1.99	8.63
II	<b>Mean</b>	6.50	47.13	96.08	51.27	30.26	42.93	2.73	3.66	123.00	16.85	8.78	160.15
	<b>± SE</b>	0.40	4.15	2.58	4.22	1.06	1.21	0.36	1.05	16.06	2.63	1.15	15.05
III	<b>Mean</b>	6.72	52.70	99.88	51.96	33.16	36.09	2.44	5.38	146.56	19.26	5.61	144.13
	<b>± SE</b>	0.52	3.83	3.30	4.85	1.84	3.15	0.16	1.64	44.27	5.72	0.92	9.97
IV	<b>Mean</b>	7.07	57.47	104.93	71.21	31.37	45.07	3.80	6.05	186.00	31.85	8.97	164.08
	<b>± SE</b>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
V	<b>Mean</b>	7.00	49.53	96.27	61.50	25.93	39.20	2.20	3.36	212.82	45.14	15.91	184.70
	<b>± SE</b>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

**Table 4. Average of inter and intra cluster distances**

Clusters	I	II	III	IV	V
I	<b>2.474</b>				
II	2.959	<b>2.180</b>			
III	3.490	3.541	<b>2.511</b>		
IV	5.649	5.899	5.762	<b>0.452</b>	
V	5.477	6.126	7.472	7.066	<b>0.000</b>

number III (5.61). Seed yield per plant revealed highest mean in cluster number V (184.70) and lowest mean in cluster number III (144.13). There was parallelism between genetic and geographical diversity. On the basis of cluster mean values for different characters it can be concluded that the varieties fall in different cluster for their respective character may be selected for hybridization programme. The viewpoint has been supported by the work of Akther et al. [8], Kumar et al. [9] and Rana et al. [10].

The average intra and inter cluster D<sub>2</sub> presented in Table 4. revealed maximum inter cluster D<sub>2</sub> value (7.472), between cluster III and V followed by cluster IV and V (7.066), whereas the minimum inter cluster D<sub>2</sub> value (2.959) was recorded between cluster I and II. The maximum intra cluster distance were found (2.511) for cluster III followed by cluster I (2.474), (2.180) for cluster II and (0.452) for cluster IV, whereas minimum intra cluster value was recorded in cluster V (0.00). "It is apparent therefore; the genotypes of cluster do not differ significantly with regards to their relative genetic distance as indicated from low variation of D<sup>2</sup> values. Crosses suggesting parent belonging to most divergence clusters would be expected to manifest maximum heterosis and also wide variability of genetic architecture. Thus, the crosses between the genetically diverse genotypes of cluster I and V based on genetic diversity and superiority with respect to any of traits the genotype may be identified and may be involve in crossing for obtaining high heterotic population, segregates and also may be exploited for development of hybrid" [4,11-18].

### 3.1 Significance of the Study

After assessing all the varieties and their characteristics it is suggested that Pusa Lal Chaulai and Pusa Kiran varieties shows better result in terms of yield attributing characters therefore these varieties can be used further in breeding improvement programmes.

## 4. CONCLUSION

The *Amaranthus* genotypes employed in this study can be successfully used for future breeding programmes, in accordance with the Mahalanobis D<sub>2</sub> analysis. For the majority of the examined traits, inter-crossing of genotypes exhibiting the greatest genetic divergence should produce superior heterotic hybrids and valued

segregants in successive generations. It is anticipated that better performing variants will be developed to boost field *Amaranthus* yield.

The individual characters contributing maximum to the D<sup>2</sup> values have greater emphasis for deciding the cluster for the purpose of further breeding improvement programme. The maximum contribution percentage was found with days to germination contributes maximum towards total divergence and this was followed by days to maturity, inflorescence length, number of leaves per plant, number of branches per plant, 50 percent flowering, fresh leaf weight, biological yield ton per hectare, seed yield kg per hectare, plant height, seed yield per plant and biological yield per plant.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Espitia-Rangel E. Breeding of grain amaranth. In: ParedesLopez O (ed.), *Amaranth: biology, chemistry and technology*. Boca Raton: CRC Press. 1994;23–38.
2. Ebert AW, Wu T, Wang S. *Vegetable amaranth (Amaranthus L.)*. AVRDC Publication no. 11-754. Tainan, Taiwan: AVRDC – The World Vegetable Center; 2011.
3. Grubben GJH. *Amaranthus L.* In: Siemonsma JS, Piluek K (Eds.), *Prosea: Plant Resources of South-East Asia 8. Vegetables*. 1993;82-86.
4. Shukla S, Bhargava A, Chatterjee A, Pandey AC, Mishra BK. Diversity in phenotypic and nutritional traits in vegetable amaranth (*Amaranthus tricolor*), a nutritionally underutilised crop. *Journal of the Science of Food and Agriculture*. 2010;90:139–144.
5. Katiyar RS, Shukla S, Rai S. Varietal performance of grain amaranths (*A. hypochondriacus*) on sodic soil. *Proceedings of the National Academy of Sciences, India, Section B: Biological Sciences*. 2000;70:185–187.
6. Schippers RR. *African indigenous vegetables: An overview of the cultivated species*. Chatham: Natural Resources Institute/ ACP-EU Technical Centre for Agricultural and Rural Cooperation; 2000.

7. Mahalanobis PC. On the generalized distance in statistics. Proceedings of National Institute of Sciences, India. 1936; 2:49–55.
8. Akther CA, Hasan M, Raihan MS, Hossain MM, Mian MAK. Genetic divergence in stem amaranth (*Amaranthus tricolor* L.) genotypes for yield and its component characters. The Agriculturists. 2013;11(1): 82-88.
9. Kumar Y. Genetic divergence analysis in *Amaranthus* (*Amaranthus* spp.) genotypes for yield and its component characters; 2019.
10. Rana JC, Yadav SK, Mandal S, Yadav S. Genetic divergence and interrelationship analysis in grain Amaranth (*Amaranthus hypochondriacus*) germplasm. Indian Journal of Genetics and Plant Breeding. 2005;65(02):99–102.
11. Arif MUHAMMAD, Jatoi SA, Rafique TARIQ, Ghafoor ABDUL. Genetic divergence in indigenous spinach genetic resources for agronomic performance and implication of multivariate analyses for future selection criteria. Journal on Science and Technology for Development. 2013;32(1):7-15.
12. Chattopadhyay A, Das S, Rana PN, Seth T, Dutta S. Estimation of genetic parameters, inter-relationships and genetic divergence of vegetable amaranths. International Journal of Plant Breeding. 2013;7(2):111-115.
13. Gerrano AS, Jansen van Rensburg WS, Adebola PO. Genetic diversity of *Amaranthus* species in South Africa. South African Journal of Plant and Soil. 2015;32(1):39-46.
14. Ngomuo M, Stoilova T, Feyissa T, Kassim N, Ndakidemi PA. The genetic diversity of leaf vegetable jute mallow (*Corchorus* spp.): A review. Indian Journal of Agricultural Research. 2017;51(5).
15. Allard RW. Principles of Plant Breeding. John Wiley and Sons, Inc. New York. 1960;485.
16. Dewey DR, Lu KH. A correlation and path coefficient analysis of components of crested wheatgrass seed production. Agronomy Journal. 1959;45:478-481.
17. Panse VG, Sukhatme PV. Statistical methods for agricultural work. 1969; 31(3):43-54.
18. Searle SR. Phenotypic, genotypic and environmental correlations. Biometrics. 1961;17:474-480.

© 2023 Tomar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/100708>