



Do Dynamometric Variables of the Pelvic Floor Muscles Differ between Women with and without Stress Urinary Incontinence? A Blind, Cross-sectional Study

Karina Moyano Amorim ^a, Elora dos Santos Silva de Lima ^a,
Caroline Razera Ferreira ^a, Alisson Monteiro de Freitas ^a,
Luis Eduardo Peixoto Rosa dos Santos ^a,
César Ferreira Amorim ^b,
Daniela Aparecida Biasotto-Gonzalez ^a
and Fabiano Politti ^{a*}

^a Physical Therapy Departament, Postgraduate Program in Rehabilitation Sciences, Universidade Nove de Julho, São Paulo, Brazil.

^b Physical Therapy Departament, Postgraduate Program in Rehabilitation Sciences, Universidade Cidade de São Paulo, São Paulo, Brazil.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMMR/2024/v36i55429

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115114>

Original Research Article

Received: 05/02/2024

Accepted: 10/04/2024

Published: 15/04/2024

*Corresponding author: E-mail: fabianopolitti@gmail.com;

ABSTRACT

Objective: The aim of the present study was to compare pelvic floor muscles (PFM) contraction variables between women with and without stress urinary incontinence (SUI).

Materials and Methods: This cross-sectional study evaluated the PFM of 17 healthy women and 17 women with SUI during a single test session using a vaginal dynamometer. Outcomes: peak time (time at which peak force occurred after the onset of contraction), passive force (baseline), maximum contraction force, impulse of contraction, average force and endurance time. Data was recorded during a single test session using a vaginal dynamometer.

Results: The following PFM contraction variables were evaluated: Analysis of covariance (ANCOVA) with the Bonferroni post hoc test was used to compare the dynamometric data between groups (control and SUI), considering age and number of childbirths as co-variables. Significant difference was observed between groups with regard to endurance ($F = 4.87, P < .03$; ANCOVA test), whereas no significant differences were found for the other variables analyzed.

Conclusion: The endurance time of PFM contraction is shorter in women with SUI, whereas variables related to the intensity of pelvic floor muscle contraction force and time from the onset to peak contraction of these muscles are similar between women with and without stress urinary incontinence.

Keywords: Physiotherapy; dynamometer; stress urinary incontinence; pelvic floor muscle.

1. INTRODUCTION

Urinary incontinence is described as any involuntary loss of urine. This condition has a tendency to aggravate over time and the prevalence figures increase with increasing age, and in women aged ≥ 70 years more than 40% of the female population is affected [1].

Stress urinary incontinence (SUI) is characterized by urinary leakage associated with some form of exertion, such as coughing, sneezing, squatting, physical exertion, and the practice of sports involving jumping, fast running, and rotational movements being, the most common form of the disorder in female population [2,3]. The cause of this phenomenon has not been fully clarified, but the loss of the integrity of the muscles that constitute the pelvic floor is one of the factors associated with this disorder [3].

In clinical practice, visual inspection and intra-vaginal palpation are the most common pelvic floor muscle (PFM) assessment methods, the latter of which is scored with different rating scales, the most often employed of which is the Oxford grading system [4,5]. These methods are minimally invasive, inexpensive and easy to perform, but are considered subjective, as the assessment of the PFM with vaginal palpation depends on the experience of the evaluator and the patient's voluntary participation as well as the positions of the patient and the examiner [6].

Moreover, divergent opinions are found in the literature regarding the reliability of these methods [5,7,8]. Thus, measurements obtained using electromyography, perineometry, dynamometry and different imaging methods (ultrasound, magnetic resonance and urodynamic analysis) have been used to determine and quantify the function of these muscles in a more precise manner [9].

Dynamometry, in particular, enables the objective evaluation of PFM force with the use of a vaginal probe. The PFM variables considered for such an evaluation include maximum strength, mean value of the strength curve, endurance, contraction velocity and passive force [10,11]. However, a study has proposed further evaluation variables based on data obtained from a vaginal dynamometer, such as impulse of contraction, average force and the peak time (time that peak force occurs after the onset of PFM contraction) [12].

The reproducibility and reliability of these new indices are high, but no comparisons have been made between women with and without SUI. Such comparisons are important, as these indices may reveal clinical characteristics (impulse of contraction, average force and the peak time) of women with SUI that have not been described in previous studies. Such information is relevant to clinical practice, as it could help guide the use of variables obtained through vaginal dynamometry as part of the diagnosis of SUI.

Therefore, the hypothesis tested in this study was that the variables passive force (baseline), maximum force, average force, impulse of contraction, time from the onset to peak contraction and endurance time have different values between women with and without SUI. Thus, the aim of the present study was to compare vaginal dynamometric variables between women with and without stress urinary incontinence during contraction of the pelvic floor muscles.

2. MATERIALS AND METHODS

2.1 Design

This was a blind, cross-sectional study conducted after being approved by the Ethics Committee of the Nove de Julho University (process n°: 1.042.129). All individuals were properly informed regarding the objectives and procedures and signed a statement of informed consent prior to testing.

2.2 Sample Size

The sample size was based on the study by Chamochumbi et al [13], considering the mean and standard deviation (\pm) of the maximum strength of PFM recorded with dynamometer (anteroposterior position) of women with and without SUI being the values respectively: 0.1 ± 0.1 N and 0.3 ± 0.2 N.

For the calculation of the sample, it was considered:

- $\alpha = 0.05$ (5% chance of a type I error)
- $1-\beta = 0.90$ (power of the sample).

A minimum of 15 individuals was determined. This calculation was performed using the G*Power 3.1.9.2 software, as recommended by Faul et al [14].

2.3 Participants

Seventeen healthy women and 17 women with SUI aged 20 to 60 years were recruited from the academic community as well as urogynecology and physical therapy services to participate in the study.

The inclusion criteria for the SUI group were a report of urinary leakage during coughing, sneezing, squatting, physical exertion, and the practice of sports involving jumping, fast running,

and rotational movements, in the previous three months and a positive answer to Question 3 of the Three Incontinence Questions used to distinguish between SUI and urgent urinary incontinence in adult women [15].

The following were the exclusion criteria: responses of "moderately" or "a lot" on the symptoms scale of the King's Health Questionnaire (KHQ) related to urgency and overactive bladder, urinary tract or vaginal canal infection, abnormal vaginal mucosa (candidiasis), current pregnancy, organ prolapse (PopQ > Phase II) [16,17], use an analgesic or muscle relaxant [10,18] and a history of urogynecology surgery.

Women without a complaint of urinary leakage and without any of the exclusion criteria were selected for the control group.

2.4 Blinding

Independent evaluators performed the following procedures: Evaluator 1: triage and evaluation of clinical characteristics; Evaluator 2: vaginal dynamometer data collection; Evaluator 3: vaginal dynamometer signal processing and statistical analysis. Evaluators 2 and 3 were blinded in relation to the groups.

2.5 Evaluation Procedures

The tests were performed in a single session. The clinical characteristics of women with SUI were assessed by a physiotherapeutic with at least 5 years of experience through the following instruments: i) the impact of urinary continence was evaluated using the International Consultation on Incontinence Questionnaire Short Form (ICIQ-SF; 0 to 21 points; slight [1–5 points], moderate [6–12 points], severe [13–18 points], and very severe [19–21 points]) [19]; ii) severity of urinary leakage was evaluated using the Protection, Amount, Frequency, Adjustment and Body Image (PRAFAB) Questionnaire (≥ 14 points: severe urinary incontinence and < 14 points: no severe urinary incontinence) [20]; and iii) quality of life was evaluated using the KHQ (0 to 100 points, with higher scores denoting poorer quality of life) [21].

The PFM contraction data were collected using a vaginal dynamometer (Model: Power Gyneco, EMG System do Brasil®) composed of a shaft measuring 70 mm in length and 22 mm in diameter and weighing 258 g. Force was

measured in Newtons (N). The dynamometer signal was acquired using a conditioner module (EMG System do Brasil®) and digitized by a 12-bit analog/digital converter with a sampling frequency of 1000 Hz connected to a computer. Both devices remained disconnected from the electrical grid during the readings to avoid interference.

The tests were performed beginning from the first day after the end of the menstrual period by a physiotherapist with experience in evaluating the PFM of women with urinary incontinence. Prior to the test, the volunteer was asked to empty her bladder and was then placed in the dorsal lithotomy position.

The volunteer was asked to perform PFM contraction prior to the test and the responses were measured by the visual inspection performed by the examiner. Training was then performed for the volunteer to learn how to contract the PFM without the contraction of other muscle groups, such as the abdominal and gluteus muscles. After these procedures, the dynamometer covered with a condom (OLLA®, Hypermarcas-S/A) was inserted into the vagina and the volunteer was instructed to relax the PFM, at which time passive force (baseline) was recorded for ten seconds. A single

command was then given for the volunteer to perform and sustain maximum contraction until exhaustion (endurance).

The data were recorded considering PFM contraction force on the sagittal (antero-posterior) plane. The insertion depth of the shaft of the dynamometer was not standardized, as the equipment was designed for the precise measurement of PFM contraction force independently of the location on the shaft at which the effort was performed [12]. The test was performed three times with a five-minute interval between trials. No volunteers reported any discomfort either during or after the test sessions.

2.6 Outcome Measures

The signal from the vaginal dynamometer was used to calculate the contraction force variables described by Nagano et al [12]: passive force (PF), impulse of contraction (IC), average force (AF), peak time (PT) and maximum strength value (MSV). Endurance was calculated from the onset of contraction to the point of exhaustion (Fig. 1). All signals were processed and analyzed using routines developed in Matlab® version 2022 (Mathworks Inc., Natick, Massachusetts, USA).




Fig. 1. Strength test of the pelvic floor muscles. Baseline: passive force. MSV: maximum pelvic floor muscles strength value; PT: peak time. IC: impulse of contraction of the muscles strength. On/Off: endurance time

2.7 Statistical Analysis

The Shapiro-Wilk test was used to determine the normality of the data distribution. Demographic data from the two groups were compared using the independent *t*-test. Analysis of covariance (ANCOVA) with the Bonferroni post hoc test was used to compare the dynamometric data between groups (control and SUI), considering age and number of childbirths as co-variables. Age and number of childbirths were entered as a covariate in these analyses because these variables were significantly different between the groups ($P < .05$). A *P*-value $< .05$ was considered indicative of statistical significance. The partial eta squared value (η_p^2) was used to calculate

the effect of the interactions, the results of which were interpreted based on Cohen (1988): < 0.01 = small effect; 0.06 = moderate effect; and ≥ 0.14 = large effect. All statistical analyses were performed with the aid of SPSS version 20.0 (IBM Corporation, Armonk, NY, USA).

3. RESULTS

Table 1 displays the demographic and clinical data of the volunteers. Age and number of childbirths differed significantly between the two groups. The PRAFAB value indicated that the group of women with SUI presented a non-severe level of urinary leakage (< 14 points) [20].

Table 1. Mean and standard deviation of demographic and clinical variables of women with (SUI) and healthy controls (women without a complaint of urinary leakage)

	SUI	Controls	P-value
Age (years)	45.35±10.61	31.24±10.23	<0.001*
Body mass index (Kg/m ²)	28.75±5.79	26.91±5.23	0.33
Parity	2.47±1.62	0.94±1.08	0.002*
PRAFAB	10.29±2.77		
ICIQ-SF	11.82±5.21		
KHQ scores			
General health perception	33.82±23.29		
Impact of incontinence	50.95±26.67		
Role limitation	39.21±28.83		
Physical limitations	46.08±35.12		
Social limitation	17.65±24.86		
Personal relationship	16.67±22.05		
Emotions	30.72±31.80		
Sleep/energy	30.39±30.75		
Severity measures	46.35±22.05		

KHQ: King's Health Questionnaire. ICIQ-SF: International Consultationon Incontinence Questionnaire Short Form.

PRAFAB: Protection, Amount, Frequency, Adjustment and Body Image Questionnaire.

* Significant difference between groups (Independent *t*-test $P < .05$)

Table 2. Mean and standard deviation values for dynamometric variables in women with stress urinary incontinence (SUI) and healthy controls (women without a complaint of urinary leakage)

	SUI	Control	F	P-value	η_p^2
Baseline (N)	6.06±1.62	6.21±1.26	0.15	.69	<0.01
MSV (N)	12.72±3.45	13.80±4.99	0.11	.91	<0.01
IC (N/s)	10.97±8.67	11.29±8.25	0.18	.66	<0.01
AF (N)	8.71±2.14	9.12±3.19	0.01	.92	<0.01
PT (s)	2.18±1.43	2.34±0.97	0.17	.67	0.01
Endurance (s)	19.14±8.15*	29.32±12.66	4.87	.03	0.14**

* Significant difference between groups (Ancova test).

** Large effect size

Baseline: passive force of the pelvic muscle floor. MSV: maximum pelvic muscle floor strength value. IC: impulse of contraction of the pelvic muscle floor. AF: Average contraction force of the pelvic muscle floor. PT: the time interval between the beginning and the peak of contraction of the pelvic floor muscles. Endurance: onset of contraction to the point of exhaustion.

The mean ICIQ-SF value of 11.82 ± 5.21 demonstrated a moderate impact (range 6–12 points) [19] of urinary incontinence on the quality of life in the women with SUI. The scores of the KHQ (quality of life; 100 points, with higher scores denoting poorer quality of life) demonstrated that the greatest impact of SUI is concentrated on urinary incontinence itself (50.95 ± 26.6) followed by physical limitations (46.08 ± 35.12), severity measures (46.35 ± 22.05) and role limitation (39.21 ± 28.8).

Table 2 displays the dynamometric findings. ANCOVA revealed a statistically significant difference between groups with regard to endurance ($F = 4.87$, $P < .03$; $\eta^2 = 0.14$), whereas no significant differences between groups were found with regard to variables related to the intensity of the PFM contraction force (PF, IC, AF and MSV) or PT.

4. DISCUSSION

The hypothesis that different vaginal dynamometric variables related to pelvic floor muscles (PFM) contraction would be different between women with and without SUI was only confirmed for endurance time, whereas no significant differences between groups were found with regard to variables related to the intensity of the PFM contraction force (PF, IC, AF and MSV) or PT.

4.1 PFM and Force

There is a growing body of evidence that lends support to the reasoning that an improvement in the tone of the PFMs can assist in improving and/or controlling SUI [22,23]. Such observations suggest the occurrence of functional alterations in the PFM of women with SUI in relation to healthy women and that components of muscle strength are associated with this disorder. Therefore, the aim of PFM training for women with SUI is generally to improve the strength, endurance and coordination of these muscles [24,25].

However, the possible alteration in the components of muscle strength in women with SUI was not confirmed by the results in the present study, as no significant differences were found for any of the variables related to the contraction force of the PFMs (PF, IC, AF and MSV) in the groups with and without SUI. Moreover, this relationship has not yet been clarified in other investigations involving

dynamometry [26,27]. Previous studies have demonstrated that the development of active strength in the tissues of the pelvic floor may [13,27] or may not [26] be significantly reduced in women with SUI. Conflicting results are also found in the analysis of passive mechanical forces measured using dynamometry with the patient at rest i.e., in one study, continent women demonstrated greater passive strength compared to incontinent women [26] while in two other studies there was no difference found between the SUI and control groups [13,27].

These different responses in the literature regarding PFM contraction force may stem from differences in the evaluation methods employed during the tests as well as differences in the equipment used to measure PFM strength. The dynamometer used in the present study was designed with a load cell positioned in the center of the shaft so that PFM contraction force could be measured precisely, independently of the location of the shaft at which the effort was performed [12].

Dynamometers used for the evaluation of the PFMs are generally constructed with strain gauges located at the base of the shaft and therefore distant from the point at which the device receives the PFM contraction force rest [13, 26,27]. This characteristic of the equipment increases the chance of a measurement error.

A previous study reports similar results regarding the association between endurance and SUI [26]. Therefore, this response indicates that the PFMs of women with SUI may be less resistant to fatigue than those of healthy women. However, this association was not found in another investigation, in which no difference was found regarding the time at which the fatigue of these muscles occurs [27]. This divergence may be explained by differences in the protocols employed for the data collection process. In the first study, before the device was inserted into the vagina each subject adopted a supine lying position with hips and knees flexed, feet flat on a conventional gynecologist's table while in the second study, the device was inserted into the vagina guided manually by the gynecologist while the women were seated semi-recumbent in a gynecological chair [27].

The time between the onset and peak PFM contraction was calculated to determine whether the time required for the recruitment of fibers in these muscles to reach maximum force is altered

in women with SUI. This possibility was not confirmed in the present study, as no difference was found between the two groups.

4.2 Implications for Physiotherapy Practice

The present findings seem to demonstrate the dynamometric variables related to PFM contraction force do not enable distinguishing between women with and without SUI and that endurance (sustained contraction) time may be the only variable that can be used for this purpose. However, it is important to stress that such observations are only valid with regard to the use of these variables as a possible way of diagnosing SUI.

In general, exercise to improve pelvic floor muscle strength, endurance, power, relaxation or a combination of these is widely used for women with stress, urgency and mixed incontinence [25]. Thus, PFM contraction force and endurance assessed by the dynamometer can be used as a tool to evaluate the effects of a training program for women with SUI.

In clinical practice, the decrease in endurance time can be a safe indication of the positive effect of a certain PFM training protocol adopted for the treatment of women with SUI. This information can assist in decision-making when managing different clinical approaches for these patients.

4.3 Research Limitations

The limitation of the present study was the fact that the volunteers were not matched for age and number of childbirths. However, the latter limitation was minimized by the inclusion of these two variables used as co-variable in the statistical analysis of the data.

5. CONCLUSIONS

In the present study, women with stress urinary incontinence demonstrated a shorter endurance time of pelvic floor muscle contraction in comparison to healthy women, whereas no differences between groups were found for variables related to the intensity of the contraction force of the pelvic floor muscles (passive force, impulse of contraction, average force and maximum strength value) or time from the onset to peak contraction of these muscles.

CONSENT

As per international standards or university standards, Participants' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

To carry out the study, the authorization of the Local Health Research Ethics Committee of the University was obtained (process nº: 1.042.129).

ACKNOWLEDGEMENTS

We thank all the patients, their carers and the health professionals who participated in the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Milsom, Gyhagen M. The prevalence of urinary incontinence. *Climacteric.* 2019;22(3):217-222.
2. De Mattos LTR, Matsuoka PK, Baracat EC, Haddad JM. Urinary incontinence in female athletes: A systematic review. *Int Urogynecol J.* 2018;29(12):1757-1763.
3. Wu JM. Stress Incontinence in Women. *N Engl J Med.* 2021;384(25):2428-2436.
4. Deegan EG, Stothers L, Kavanagh A, Macnab AJ. Quantification of pelvic floor muscle strength in female urinary incontinence: A systematic review and comparison of contemporary methodologies. *Neurorol Urodyn.* 2018;37(1):33-45.
5. Pena CC, Bø K, de la Ossa AMP, Fernandes ACNL, Aleixo DN, de Oliveira FMF, Ferreira CHJ. Are visual inspection and digital palpation reliable methods to assess ability to perform a pelvic floor muscle contraction? An intra-rater study. *Neurorol Urodyn.* 2021 Feb;40(2):680-687.
6. Bø K, Finckenhagen HB. Vaginal palpation of pelvic floor muscle strength: inter-test reproducibility and comparison between palpation and vaginal squeeze pressure. *Acta Obstet Gynecol Scand.* 2001;80:883-887.
7. da Silva JB, de Oliveira Sato T, Rocha APR, Driusso P. Inter- and intrarater

- reliability of unidigital and bidigital vaginal palpation to evaluation of maximal voluntary contraction of pelvic floor muscles considering risk factors and dysfunctions. *Neurourol Urodyn.* 2021 Jan;40(1):348-357.
8. da Silva JB, de Godoi Fernandes JG, Caracciolo BR, Zanello SC, de Oliveira Sato T, Driusso P. Reliability of the PERFECT scheme assessed by unidigital and bidigital vaginal palpation. *Int Urogynecol J.* 2021 Dec;32(12):3199-3207.
 9. Bo K, Frawley HC, Haylen BT, et al. An International Urogynecological Association (IUGA)/International Continence Society (ICS) joint report on the terminology for the conservative and nonpharmacological management of female pelvic floor dysfunction. *Int Urogynecol J.* 2017;28(2):191-213.
 10. Dumoulin C, Bourbonnais D, Lemieux MC. Development of a dynamometer for measuring the isometric force of the pelvic floor musculature. *Neurourol Urodyn.* 2003;22:648-653.
 11. Martinho NM, Marques J, Silva VR, Silva SLA, Carvalho LC, Botelho S. Intra and inter-rater reliability study of pelvic floor muscle dynamometric measurements. *Braz J Phys Ther.* 2015;19:97-104.
 12. Nagano RCR, Biasotto-Gonzalez DA, da Costa GL, Politte F. Test-retest reliability of the different dynamometric variables used to evaluate pelvic floor musculature during the menstrual cycle. *Neurourol Urodyn.* 2018;17:1.
 13. Chamochumbi CCM, Nunes FR, Guirro RRJ, Guirro ECO. Comparison of active and passive forces of the pelvic floor muscles in women with and without stress urinary incontinence. *Braz J Phys Ther.* 2012;16:314-319.
 14. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods.* 2007;39:175-191.
 15. Brown JS, Bradley CS, Subak LL, et al. The Sensitivity and Specificity of a Simple Test To Distinguish between Urge and Stress Urinary Incontinence. *Ann Intern Med.* 2006;144:715-723.
 16. Brusaca LA, Rocha APR, Cardozo L, Oliveira AB, Driusso P. Brazilian version of the King's Health Questionnaire: Assessment of the structural validity and internal consistency in female urinary incontinence. *Int Urogynecol J.* 2022;33(11):3143-3154.
 17. Dumoulin C, Bourbonnais D, Lemieux MC. Development of a dynamometer for measuring the isometric force of the pelvic floor musculature. *Neurourol Urodyn.* 2003;22:648-653.
 18. Hundley AF, Wu JM, Visco AG. A comparison of perineometer to brink score for assessment of pelvic floor muscle strength. *Am J Obstet Gynecol.* 2005;192:1583-1591.
 19. Kloving A, Avery K, Sandvik H, Hunskaar S. Comparison of two questionnaires for assessing the severity of urinary incontinence: The ICIQ-UI SF versus the incontinence severity index. *Neurourol Urodyn.* 2009;28(5):411-5.
 20. Hendriks EJ, Bernards AT, Staal JB, de Vet HC, de Bie RA. Factorial validity and internal consistency of the PRAFAB questionnaire in women with stress urinary incontinence. *BMC Urol.* 2008;8:1.
 21. Tamanini JTN, D'Ancona CAL, Botega NJ, Rodrigues Netto N. Validation of the Portuguese version of the King's Health Questionnaire for urinary incontinent women. *Rev Saude Publica.* 2003;37:203-211.
 22. Luginbuehl H, Baeyens JP, Taeymans J, Maeder IM, Kuhn A, Radlinger L. Pelvic floor muscle activation and strength components influencing female urinary continence and stress incontinence: a systematic review. *Neurourol Urodyn.* 2015;34:498-506.
 23. Price N, Dawood R, Jackson SR. Pelvic floor exercise for urinary incontinence: A systematic literature review. *Maturitas.* 2010;67:309-315.
 24. Alves FK, Riccetto C, Adami DB, Marques J, Pereira LC, Pama P, et al. A pelvic floor muscle training program in postmenopausal women: A randomised controlled trial. *Maturitas.* 2015;81(2):300-5.
 25. Dumoulin C, Cacciari LP, Hay-Smith EJ. Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women. *Cochrane Database of Systematic Reviews.* 2018, Issue 10. Art. No: CD005654.

26. Morin M, Bourbonnais D, Gravel D, Dumoulin C, Lemieux MC. Pelvic floor muscle function in continent and stress urinary incontinent women using dynamometric measurements. *Neurorol Urodyn*. 2004;23:668-674.
27. Verelst M, Leivseth G. Force and stiffness of the pelvic floor as function of muscle length: A comparison between women with and without stress urinary incontinence. *Neurorol Urodyn*. 2007;26: 852-857.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/115114>*