



Evaluation of Probiotic Attributes of Isolated Yeast Strains from Fruits and Bread

Malashree L^{a++}, Anushree Y K^{a++} and Akshaykumar^{b++*}

^a Dairy Science College, Hebbal, Bangalore, India.

^b Dairy Science College, Mahagaon Cross, Kalaburagi, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ejnf/2024/v16i81499>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118821>

Original Research Article

Received: 15/05/2024

Accepted: 17/07/2024

Published: 25/07/2024

ABSTRACT

Aims: To characterize the probiotic attributes of yeasts obtained from fruit and bread samples.

Study Design: Isolation, characterization and performance evaluation for probiotic attributes.

Place and Duration of Study: Department of Dairy Microbiology, Dairy Science College, Hebbal, Bengaluru, Karnataka Veterinary, Animal and Fisheries Sciences University, Bidar, Karnataka, India, between June 2022 and March 2023.

Methodology: 3 samples (Litchi, orange and bread) were taken to isolate yeasts using pour plate method. The obtained colonies were isolated, purified and cultured on YMM (Yeast maintenance media) and checked for preliminary identification, biochemical tests followed by probiotic characterization using acid and bile test.

⁺⁺ Assistant Professor;

*Corresponding author: Email: akshaysagar200@gmail.com;

Cite as: L, Malashree, Anushree Y K, and Akshaykumar. 2024. "Evaluation of Probiotic Attributes of Isolated Yeast Strains from Fruits and Bread". *European Journal of Nutrition & Food Safety* 16 (8):110-16. <https://doi.org/10.9734/ejnf/2024/v16i81499>.

Results: The yeast counts were only obtained from litchi and orange but not in bread sample. The counts were 4.20, 3.3 and 0 log₁₀cfu/g for litchi, orange and bread respectively with significant difference ($P=0.05$). Isolated yeasts of 5 numbers were identified as *Saccharomyces* species and those were subjected for acid and bile tolerance test to check probiotic attributes. All the isolates showed tolerance to both acid and bile with significant difference.

Conclusion: The isolated yeast strain possess some probiotic characteristics and use of probiotic yeasts by humans through food gives biotherapeutic value.

Keywords: Probiotics; acid and bile tolerance; saccharomyces; biotherapeutic.

1. INTRODUCTION

Yeast constitutes a diverse group of single-celled fungi, playing pivotal roles in various biological processes. Classified under the kingdom Fungi, yeast species are widespread in nature, thriving in environments ranging from soil and water to the surfaces of plants and animals [1]. *Saccharomyces cerevisiae* is one of the most well-known yeast species and is extensively used in food and beverage fermentation. The characteristics of yeast encompass a broad array of features that define its biology and behavior. One notable trait is its reproductive versatility [2].

Yeast cells can reproduce both sexually and asexually, providing them with flexibility in different environments. Asexual reproduction often occurs through budding, where a smaller cell (bud) forms on the parent cell, eventually detaching to become an independent organism. Genetic plasticity is another significant characteristic [3].

Different yeast species exhibit preferences for specific temperature ranges, impacting their performance in various environments. Most yeast strains thrive in a moderate temperature range, approximately 25-30 °C. This mesophilic preference aligns with the typical conditions found in many ecological niches, including fermenting fruits and the human body. The appearance of yeast is a diverse and fascinating aspect that varies among species and can be observed at both macroscopic and microscopic levels. Under a microscope, typical yeast cells exhibit distinctive shapes, commonly appearing as ovals or spheres.

Probiotics, often associated with bacteria, also include certain types of yeast that confer health benefits when consumed in adequate amounts. *Saccharomyces boulardii* is a prominent example of a probiotic yeast known for its therapeutic properties. This yeast has gained attention for its

potential positive impact on gastrointestinal health and overall well-being [4].

Gut microbiota is a complex community of microorganisms that live in the digestive tracts of humans and animals, including insects. In people, the gut microbiota has the biggest quantities of microorganisms, and the greatest number of species compared to other parts of the body [5]. One of the primary role of probiotic yeast is in maintaining a balanced microbial community in the gut. The human gastrointestinal tract is home to a complex ecosystem of microorganisms, collectively known as the gut microbiota. This microbiota plays a crucial role in digestion, nutrient absorption, and the overall health of the digestive system. Probiotics, including certain yeast strains, contribute to the diversity and balance of this microbial community [6].

2. MATERIALS AND METHODS

All the experiments were conducted at the post graduate laboratory of Dairy Microbiology Department, Dairy Science College, Hebbal, Bengaluru, Karnataka Veterinary, Animal and Fisheries Sciences University, Regional Campus.

2.1 Materials

2.1.1 Fruit and bread samples

Litchi, orange and bread samples were purchased from the local market, Hebbal, Bangalore, Karnataka for isolating yeast.

2.1.2 Yeast Maintenance Media (YMM)

Media used for isolating yeast from sample.

2.1.3 Physiological saline

Diluent used for serial dilution of samples.

2.1.4. For biochemical tests

Hugh and Leifson's medium, Carbohydrate fermentation broth, Christensen's Urea Agar, Nitrate broth is used.

2.2 Methods

2.2.1 Enumeration and characterization of yeast from fruits and bread

2.2.1.1 Enumeration of yeast

Enumeration was performed by using plate count method. From the collected orange and litchi samples, peel the outer layer of the fruit and pieces of bread sample were transferred to the prepared yeast maintenance media (YMM broth) of 11g and leave it for enumeration overnight. On next day serial dilution was performed using physiological saline serially like 1:10, 1:100, 1:1000. 1ml of each serially diluted samples were transferred to sterile petri plates and 10ml of sterile molten YMM agar were transferred and kept for incubation at 30°C for 3-5 days. After the completion of the incubation period, the colonies on YMM agar were counted. The countable plates ranging between 30- 300 were counted and average count was expressed as log₁₀cfu/g. From selected countable plates, colonies were selected, colony morphology noted and purified by streaking on to YMM agar plates three times and maintained on YMM agar slants. Once in a month, the isolates were sub cultured and maintained in refrigerator at 5 °C.

2.2.1.2 Characterization of obtained isolates

Purified isolates were subjected for preliminary test to confirm the genus and specific tests to speciate.

2.2.1.2.1 Preliminary identification of isolates

Maintained isolates were subjected for preliminary test like simple staining as per the standard procedures of Harrigan [7].

2.2.1.2.2 Specific tests for the isolates

After confirming the status of isolates through the preliminary tests, all the isolates were made to carry out oxidative and fermentative test, Urease test, Nitrate reduction test, carbohydrate fermentation test as per the standard procedures of Harrigan [7].

2.2.1.2.3 1 of (Oxidative- Fermentative) test

Prepared 2 ml of HL medium in test tubes was sterilized by autoclaving at 121 °C for 15 min. Inoculated the test culture into HL medium tubes by stabbing. To one of the tubes layer sterile paraffin wax as a sealant to cover ½ to 1 cm. Incubate the tubes at 37 °C for 48 h to 72 hrs. change in colour from blue to yellow indicates acid production and positive for test. Blue as such or darkened blue indicate no acid production thus negative for the test.

2.2.1.2.4 Urea hydrolysis

Actively grown young broth culture was inoculated on the slope part of Christensen's urea agar slants and incubated at 37 °C for 24-48 h. Urease production and subsequent hydrolysis of urea resulted in the production of ammonia which increased the pH and this indicated by a change in colour of the medium from yellow to pink was considered as positive.

2.2.1.2.5 Nitrate reduction test

Inoculate the glucose nitrate broth with the fresh culture and incubate at 37°C for 24 to 48 hrs. Add 1 ml of each test reagent (Griess-Ilosvay's reagent) to the test culture and look for colour change. A red colour development within few minutes indicates the presence of nitrite and hence the ability of the microorganisms to reduce nitrate to nitrite.

2.2.1.2.6 Carbohydrate fermentation test

Prepare Carbohydrate fermentation broth and transfer 10ml quantity in sterilize test tube, sterilize by autoclaving. Prepare the sugars (Glucose and Lactose) at 10% concentration each and filter sterilize. Transfer 1 ml of sterile sugar to 10 ml of broth □ Inoculate the culture and incubate, observe for colour which has pH indicator with gas production in Durham's tube. Change in colour from purple to yellow with turbidity indicates positive for the test with homofermentation. In heterofermentation gas production is observed in Durham's tube. No change in colour indicates negative for test.

2.2.2 Evaluation of probiotic attributes of yeast isolates

2.2.2.1 Acid tolerance test

With slight modification, the method of Dixon et al [8] YMM broth was prepared in 3 different test

tubes with 3 different pH like 1.0, 2.0, 3.0. Yeast isolates are inoculated into 3 test tubes, after inoculation the culture were inoculated onto the YMM agar plates and checked for the growth of an organism which is incubated at 30°C for 72 hours.

2.2.2.2 Bile tolerance test

For testing the bile tolerance, YMM broth is taken into 3 test tubes. To this ox-bile is prepared and added in a different concentration like 0.1, 0.2, 0.3% [9]. Active cultures of yeast isolates are inoculated and the culture are immediately inoculated onto the YMM agar plates and checked for the growth of an organism which is incubated at 30°C for 72 hours.

3. RESULTS AND DISCUSSION

3.1 Enumeration and Characterization of Yeast from Fruits and Bread

Samples when plated for enumeration on selective media for yeast counts were ranged from 0 to 4.2 log₁₀ cfu/g. Among the 3 samples litchi showed highest count of 4.2 log₁₀ cfu/g followed by orange of 3.3 log₁₀ cfu/g and bread sample showed no yeast counts. A total of 5

isolates (Table 1) were obtained, among that 3 isolates from litchi and 2 isolates from orange were selected from colonies of countable plates (Fig.1).

Preliminary identification of the yeast isolates was revealed through simple staining. Smears of yeast isolates are prepared from the slant cultures and are heat fixed and stained with borax methylene and observed under 100x. The following biochemical tests were performed to characterize obtained isolates by subjecting to oxidative fermentative test, which showed a positive result with color change from bluish green to yellow colour with growth was seen in the HL (Hugh and Leifson's) media followed by urease test displayed a positive result on christensen urease slant agar, i.e., red colour slant changed to pink colour slant. For nitrate reduction test it showed positive with change in colour, which indicated nitrate has been reduced to nitrogen then sugar fermentation test (Sucrose, Fructose, Lactose, and Glucose were the four sugars used). Sucrose, fructose and glucose showed positive results with acid and gas production, whereas lactose showed negative results, and it did not show acid or gas production. Based on these results obtained it showed that yeast isolates were belonged to Saccharomyces family.

Table 1. Colony and cell morphology of yeast isolates obtained from fruit and bread samples

Sources of isolates	Code of isolates	Number of isolates	Colony morphology	Cell morphology
Litchi	Y1, Y2, Y3	3	Circular shape, creamish white in colour, raised elevation, shiny surface	Oval shaped cells with budding
Orange	Y4, Y5	2		
Bread	--	0		



Fig. 1. Colonies of yeast on YMM agar

In accordance with the study done by Chandimala et al [10], on isolation and characterization of yeast from locally available foods like fruits and vegetables. Yeasts were isolated and enumerated using pour plate technique. Yeast Potato Dextrose (YPD) medium was used for growing yeast and incubation was done for 1-2 days at 25°C. The discrete isolated colonies were purified by re-streaking on YPD plates and maintained on slants with 2% agar of the same medium at 5°C in the refrigerator and as frozen stocks in liquid YPD medium with 40% glycerol at -20°C. A total of 24 isolates were obtained and those were characterized by some biochemical tests like catalase test, urease test, sugar fermentation test.

3.2 Screening of Yeast Isolates for Probiotic Properties

3.2.1 Screening of yeast isolates for acid tolerance

Sterile pre-adjusted YMM broth to pH 1.0, 2.0 and 3.0 were inoculated individually with 5 yeast isolates and incubated at 30 °C. Samples drawn immediately after inoculation considered as 0 h and after 72 h of incubation were plated. Isolates at 0hr showed viable count ranged between 4.04 to 4.18 log₁₀ cfu/ml at pH 1.0, at pH 2.0 the viable count ranges between 4.00 to 4.36 log₁₀ cfu/ml and at pH 3.0 the viable count ranges between 4.20 to 4.53 log₁₀ cfu/ml whereas the isolates after 72 h ranged with respect to viable count between 5.38 to 5.74 log₁₀ cfu/ml at pH 1.0, at pH 2.0 the viable count ranges between 4.54 to 5.39 log₁₀ cfu/ml and at pH 3.0 the viable count ranges between 4.30 to 5.17 log₁₀ cfu/ml on sterile YMM agar medium.

All the isolates showed growth in different pH of 1.0, 2.0 and 3.0. As the pH increases the viability of each isolate get decreased.

3.2.2 Screening of yeast isolates for bile tolerance

The bile salt tolerance of all the 5 yeast isolates were tested in the presence of 0.1, 0.2 and 0.3 per cent ox bile. Samples drawn immediately after inoculation (0 h) and after 72h h of incubation at 30 °C were plated using sterile YMM agar. Isolates at 0 h ranged in viable count between 3.3 to 4.39 log₁₀ cfu/ml for 0.1%, 3.9 to 4.47 log₁₀ cfu/ml for 0.2% and for 0.3% at 0hr ranges between 4.07 to 4.64log₁₀ cfu/ml, whereas at 72 hr the viable count ranges between 4.65 to 5.47 log₁₀ cfu/ml for 0.1%, 4.34 to 5.44 log₁₀ cfu/ml for 0.25 and for 0.3% the viable count at 72h ranges between 4.14 to 5.23 log₁₀ cfu/ml.

All the isolates showed growth in different concentration of ox-bile. As the bile concentration increases the viability of each isolate get decreased.

Sunita et al [11,12] reported acid tolerance property for the six yeast isolates by incubating them at four different pH values i.e. 2, 2.5, 3 and 7 for different time intervals (1,3 and 5 hours). It was observed that out of all tested isolates, 1 isolate was highly tolerant to all the acidic pH values tested. 2 isolates showed good tolerance to acidic pH of 2 and 3 when incubated for different duration at these pH values. The other 3 isolates were found to be susceptible to acidic pH. All the isolates showed decrease in growth at pH 7. For testing tolerance to bile salts, the yeast

Table 2. Acid tolerance of yeast isolates obtained from fruit and bread samples

Isolate	Acid tolerance viable count (log ₁₀ cfu/ml)					
	pH 1		pH 2		pH3	
	0h	72h	0h	72h	0h	72h
Y1	4.04 ^a	5.38 ^a	4.27 ^{ab}	5.25 ^a	4.44 ^{ab}	4.97 ^a
Y2	4.17 ^a	5.47 ^a	4.36 ^a	5.39 ^a	4.53 ^a	5.17 ^a
Y3	3.40 ^b	4.74 ^b	4.00 ^b	4.54 ^b	4.20 ^b	4.30 ^b
Y4	4.18 ^a	5.42 ^a	4.25 ^{ab}	5.32 ^a	4.51 ^a	5.00 ^a
Y5	4.17 ^a	5.47 ^a	4.36 ^a	5.39 ^a	4.53 ^a	5.17 ^a
CD (P=.05)	0.18	0.15	0.22	0.14	0.16	0.14

Note:

- Values are average of three trails
- CD - Critical difference
- Higher value in the table was compared with other values
- Same superscripts within the column indicate no significant difference
- Different superscripts within the column are compared with other values

Table 3. Bile tolerance of yeast isolates obtained from fruit and bread samples

Isolate	Bile tolerance viable count (\log_{10} cfu/ml)					
	0.1%		0.2%		0.3%	
	0h	72h	0h	72h	0h	72h
Y1	4.14 ^a	5.28 ^a	4.33 ^a	5.04 ^b	4.47 ^a	4.91 ^b
Y2	4.39 ^a	5.47 ^a	4.47 ^a	5.44 ^a	4.64 ^a	5.23 ^a
Y3	3.30 ^b	4.65 ^b	3.90 ^b	4.34 ^c	4.07 ^b	4.14 ^c
Y4	4.33 ^a	5.45 ^a	4.37 ^a	5.44 ^a	4.50 ^a	5.22 ^a
Y5	4.39 ^a	5.47 ^a	4.47 ^a	5.44 ^a	4.64 ^a	5.23 ^a
CD ($P=0.05$)	0.19	0.14	0.12	0.12	0.17	0.10

Note:

- Values are average of three trails
- CD - Critical difference
- Higher value in the table was compared with other values
- Same superscripts within the column indicate no significant difference
- Different superscripts within the column are compared with other values

isolates were incubated in the presence of 5% and 10% of bile salts and their tolerance was recorded in terms of their absorbance values. Out of all the six isolates tested, 1 isolate showed maximum tolerance to bile salt followed by next 3 isolates. The other two isolates seem to be susceptible to the presence of bile salts [13,14].

4. CONCLUSION

Fermented foods are a source of diverse microflora and the fact that very few types of yeast have been commercially used as probiotic, it is worthwhile to explore the diversity of yeasts in traditional Indian fermented foods and study their probiotic attributes. Through this research we have isolated yeasts with potential probiotic properties. They could be prescribed to patients on prolonged antibiotic treatment, protect against food borne enteric pathogens. They have the ability to degrade anti-nutrients such as phytic acid and tannic acid, thereby, improving the nutritional status of food. They can hydrolyze lactose, thus prove beneficial to lactose intolerant patients. They also produce L-asparaginase which has anticancer property. These yeast isolates could aid in digestion because of their ability to produce lipase and protease. Enhance vitamin B12 pool due to its production by these isolates. They can be used as immune-stimulants. Their ability to assimilate cholesterol could be of immense value to patients suffering from hypercholesterolemia. Beneficial attributes of *Saccharomyces cerevisiae*, *Candida sp.*, *Pichia sp.*, and *Aureobasidium sp.* indicates them to be promising probiotic agents in future and can be widely used as food and feed supplements.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENTS

We would like to thank the Department of Dairy Microbiology, Dairy Science College, Hebbal, Bangalore, Karnataka, India for financial assistance for this investigation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Borkar SG, Shinde K. Yeast species of diverse functionality in health sciences: A concise report. GSC Biological and Pharmaceutical Sciences. 2023;25(2):149-168.
2. Wallen RM, Perlin MH. An overview of the function and maintenance of sexual reproduction in dikaryotic fungi. Frontiers in Microbiology. 2018;9:3243
3. Ansari F, Alian Samakkhah S, Bahadori A, Jafari SM, Ziaee M, Khodayari MT, Pourjafar H. Health-promoting properties of *Saccharomyces cerevisiae* var. *boulardii* as a probiotic; Characteristics, isolation, and applications in dairy products. Critical Reviews in Food Science and Nutrition, 2023;63(4):457-485.

4. Fakruddin MD, Hossain MN, Ahmed MM. Antimicrobial and antioxidant activities of *Saccharomyces cerevisiae* IFST062013, a potential probiotic. *BMC Complementary and Alternative Medicine*. 2017;17:1-11.
5. Gomaa E.Z. Human gut microbiota/microbiome in health and diseases: A review. *Antonie van Leeuwenhoek*. 2020;113. Available:<https://doi.org/10.1007/s10482-020-01474-7>
6. Roto SM, Rubinelli PM, Ricke SC. An introduction to the avian gut microbiota and the effects of yeast-based prebiotic-type compounds as potential feed additives. *Front. Vet. Sci*. 2015;2:28. DOI: 10.3389/fvets.2015.00028
7. Harrigan F, Wilkie. *Laboratory methods in food and dairy microbiology*. III edition, Academic press Inc. (London) Ltd., U.K; 1998.
8. Dixon B., Kilonzo-Nthenge A, Nzomo M, Bhogju S, Nahashon, S. Evaluation of selected bacteria and yeast for probiotic potential in poultry production. *Microorganisms*. 2022;10(4):676.
9. Ragavan ML, Das N. Isolation and characterization of potential probiotic yeasts from different sources. *Asian J Pharm Clin Res*. 2017;10(4):451-45.
10. Chandimala UR, Rajawardhana DU, Liyanage PLN, Hewajulige IGN. Isolation and characterization of yeasts from locally available foods. *Journal of Agro-Technology and Rural Sciences*. 2022; 2(2):1-12.
11. Sunita A, Aparajita S, Aayushi R, Arti N. Evaluation of the probiotic potential of yeasts isolated from Indian fermented food items. *Research Journal of Biotechnology*. 2021;16:10.
12. Uzoh CV, Orji JO, Okeh CO, Nworie CO, Igwe PC, Uwanta LI. Assessment of probiotic potentials of lactic acid bacteria isolated from some locally fermented foods. *Asian Journal of Food Research and Nutrition*. 2022;1(1):11–16. Available:<https://journalajfrn.com/index.php/AJFRN/article/view/2>
13. Vandana KV, Harinivenugopal Shashi Kumar CS, Malashree L, Shilpashree BG, Ramachandra B. *Bacillus Spp.* – A Potential Probiotic. *Journal of Scientific Research and Reports*. 2024;30(6):646-52. Available:<https://doi.org/10.9734/jsrr/2024/v30i62082>
14. Amorim JC, Piccoli RH, Duarte WF. Probiotic potential of yeasts isolated from pineapple and their use in the elaboration of potentially functional fermented beverages. *Food Research International*. 2018;107:518-27.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/118821>