



The Frequency of Prescribing Nicotine and Varenicline in a Public Hospital

Nehad J. Ahmed ^{a*}, Gamal A. Gabr ^b and Abeer A. El-Sherbiny ^c

^a Department of Clinical Pharmacy, College of Pharmacy, Prince Sattam Bin Abdulaziz University, Al-Kharj, Saudi Arabia.

^b Department of Pharmacology, College of Pharmacy, Prince Sattam Bin Abdulaziz University, Al-Kharj, Saudi Arabia.

^c Department of Medical Laboratory Sciences, College of Applied Medical Sciences, Prince Sattam Bin Abdulaziz University, Al-kharj, Saudi Arabia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i55A33809

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/77737>

Original Research Article

Received 01 October 2021
Accepted 02 December 2021
Published 13 December 2021

ABSTRACT

Aim: This study was conducted to demonstrate the prescribing pattern of nicotine and varenicline in a public hospital in Alkharj.

Methodology: This is a retrospective study includes evaluating outpatient prescribing of nicotine and varenicline from 1st of January 2018 to 30th of June 2018 in a public hospital in Alkharj.

Results: All of the nicotine patch prescriptions were written by internal medicine department and all of the prescriptions were prescribed by residents. The age of more than half of them was between 30 and 39 (53.33%). All of the varenicline tablet prescriptions were written by internal medicine department. The age of about 88.23% of the patients who received varenicline tablet was more than 29 years.

Conclusion: Nicotine and varenicline prescribing was infrequent in Al-kharj. More awareness programs are needed for health care workers and for the public. Moreover, the physicians should assess if the treatment is needed or no according to the person condition.

Keywords: Nicotine; outpatient; pattern; prescribing; varenicline.

1. INTRODUCTION

Cigarette smoking is a major modifiable health risk factor in the United States, significantly contributing to deaths from cancer and pulmonary and cardiovascular diseases [1]. Although it is estimated that smoking-related illnesses lead to 443,000 premature deaths and almost \$100 billion in lost productivity yearly [2]. Among, American adults, 22 % of men and 17.5 % of women still smokes regularly [3]. Among Saudis, about 12.1% of Saudis reported that they currently smoke tobacco, most of them are males (23.7% of males and 1.5% of females in Saudi Arabia smoke tobacco) [4].

Tobacco use can lead to tobacco dependence and several serious health problems. Quitting smoking significantly decreases the risk of developing smoking-related diseases [5]. In 2015, 68.0% of adult smokers in the United States (22.7 million) said that they wanted to quit smoking [6]. In 2018, 55.1% of adult smokers in the United States (21.5 million) said that they had made a quit attempt in the past year [7].

If the patients cannot quit smoking, they should be offered pharmacologic assistance in quitting, such as nicotine replacement therapies, varenicline, and bupropion. Use of these therapies during smoking cessation can double the rate of successful abstinence. In some patients, it is important to use more than one type of nicotine replacement therapy and combining these therapies with other treatments such as bupropion to provide additional benefit [1].

Nicotine is used to help treat addiction to or dependence on smoking cigarettes. Quitting smoking abruptly can lead to experience numerous severe effects and cravings that are called withdrawal symptoms [8]. So, products that deliver low doses of nicotine could be used to ease the quitting process and manage withdrawal symptoms [8].

Nicotine replacement can come in different forms including patches, lozenges, gum, inhalers, and nasal sprays [9]. It may cause several side effects on both the body and mind, including heightened mood, decreased appetite, increased activity in the intestines, increased heart rate, increased production of saliva and phlegm, sweating, increased blood pressure, diarrhea, nausea, increased alertness, better memory [10].

Varenicline is in a class of medicines called smoking cessation aids. It works by blocking the pleasant effects of nicotine on the brain [11]. It is the first approved nicotinic receptor partial agonist [12]. To increase the chance of success, this medicine should be used with a stop-smoking program that includes support, education, and counseling [13]. Varenicline can cause several adverse Effects to the patients. The most common adverse effects include nausea, abnormal dreams, headache, insomnia, in addition to several other less common effects [14].

It is important also to explore the pattern of using smoking cessation medications to ensure that they are used correctly. So, this study was conducted to demonstrate the prescribing pattern of nicotine and varenicline in a public hospital in Alkharj.

2. METHODOLOGY

This is a retrospective study includes evaluating outpatient prescribing of nicotine and varenicline from 1st of January 2018 to 30th of June 2018 in a public hospital in Alkharj that is a city in Saudi Arabia that include about 425,300 persons.

The inclusion criteria include the prescriptions that contain nicotine and varenicline during the study period and the exclusion criteria include the prescriptions before 1st of January 2018 or after 30th of June 2018 and the prescriptions that didn't contain nicotine and varenicline.

The data include the age of the patients who received nicotine, the age of the patients who received varenicline, the prescribed dosage forms, the number of prescriptions that contain nicotine or varenicline during different months, and the level of prescribers who prescribed nicotine and varenicline. The data were collected and analyzed using excel sheet and the descriptive data were represented as numbers and percentages.

3. RESULTS AND DISCUSSION

Smoking cessation medications were prescribed for 32 patients during the study. About 46.87% of them received nicotine patch and 53.13% of them received varenicline tablet. Table 1 show the dosage form of the smoking cessation

medications that were prescribed during the study period.

All of the nicotine patch prescriptions were written by internal medicine department and all of the prescriptions were prescribed by residents. The age of more than half of them was between 30 and 39 (53.33%). Table 2 shows the personal data of the patients who received nicotine patch.

Table 3 shows the number of nicotine prescriptions that were prescribed during different months of the study. More than half of the prescriptions were prescribed in April (53.33%).

Table 4 shows the age of the patients who received varenicline tablet. The age of about 88.23% of the patients who received varenicline tablet was more than 29 years.

All of the varenicline tablet prescriptions were written by internal medicine department. About 35.29% of the prescriptions were prescribed in February. The number of varenicline tablet prescriptions that were prescribed during different months of the study is shown in Table 5.

Table 6 shows the level of the prescribers who prescribed varenicline tablet. About 94.12% of the prescriptions were written by residents.

The use of smoking cessation medications was uncommon in Al-kharj. Fiore et al stated that the existence of treatment guidelines recommends the use of a smoking cessation medication for all smokers [15]. Nonetheless, only 2.4% of clinician visits by smokers result in a prescription for a cessation medicine [16]. Babb et al informed that despite the availability of evidence-based treatments to achieve smoking cessation, less than one-third of adult cigarette smokers who try to quit use any type of cessation counseling and/or FDA-approved cessation medication [6]. Previous studies also reported that most smokers attempt to quit without using treatment, with success rates of about 7–8% [6,15,17,18]. There are several caused for the uncommon use such as the common lack of health insurance among tobacco users [19]; inadequate private and public insurance coverage of cessation treatments [20]; insufficient and cumbersome reimbursement for cessation treatments offered by physicians and hospitals [21]; insufficient promotion of cessation treatments to smokers and healthcare providers, which can contribute to low use of these treatments [21]; underfunding of state tobacco quitlines and other cessation services [22,23]; and inadequate integration of tobacco use screening and cessation interventions into routine clinical care [6].

Table 1. The dosage form of the smoking cessation medications

Dosage form	Number	Percentage
Nicotine patch	15	46.87
Varenicline tablet	17	53.13
Total	32	100.00

Table 2. The Age of the patients who received nicotine patch

Variable	Category	Number	Percentage
Age	20-29	3	20.00
	30-39	8	53.33
	40-49	3	20.00
	More than 49	1	6.67

Table 3. The number of nicotine prescriptions that were prescribed during different months

Month	Number	Percentage
January	1	6.67
February	2	13.33
March	0	0.00
April	8	53.33
May	4	26.66
June	0	0.00

Table 4. The age of the patients

Variable	Category	Number	Percentage
Age	Less than 20	1	5.88
	20-29	1	5.88
	30-39	5	29.41
	40-49	5	29.41
	More than 49	5	29.41

Table 5. The number of varenicline tablet prescriptions that were prescribed during different months

Month	Number	Percentage
January	4	23.53
February	6	35.29
March	1	5.88
April	4	23.53
May	1	5.88
June	1	5.88

Table 6. The level of the prescribers who prescribed varenicline tablet

Prescriber level	Number	Percentage
Consultant	1	5.88
Specialist	0	0.00
Resident	16	94.12

Smoking cessation medications should be used only for some smokers because its use could cause several side effects. Ebbert et al reported that smoking cessation medications were well tolerated with comparable adverse events profiles and that most adverse events are not clinically important, and prescribers can reassure patients that those experienced will be manageable [24]. Signs of severe nicotine dependence in people who smoke include smoking more than 1 pack a day, smoking even while sick, smoking within 5 minutes of waking up, smoking to ease symptoms of withdrawal, and waking up at night to smoke [25].

Smoking cessation aids are available over the counter and on prescription in the several forms including gum, lozenges, tablets, inhalants, sprays, and skin patches [26]. About 46.87% of them received nicotine patch and 53.13% of them received varenicline tablet. Varenicline is available as tablets only but nicotine available as inhaler, sprays, patches and other forms but it is prescribed as patches only.

4. CONCLUSION

Nicotine and varenicline prescribing was infrequent in Al-kharj due to several causes such as the lack of awareness and insufficient

promotion of cessation treatments to smokers and healthcare providers. More awareness programs are needed for health care workers and for the public. Moreover, the physicians should assess if the treatment is needed or no according to the person condition.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT

It is not applicable.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

ACKNOWLEDGEMENT

This Publication was supported by the Deanship of Scientific Research at Prince Sattam bin Abdulaziz University.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. AAFP. Promoting Smoking Cessation. Access 13 November 2021. Available: <https://www.aafp.org/afp/2012/0315/p591.html>.
2. Centers for Disease Control and Prevention. Smoking-attributable mortality, years of potential life lost, and productivity losses—United States, 2000–2004. *MMWR Morb Mortal Wkly Rep.* 2008;57(45):1226–1228.
3. Pleis JR, Ward BW, Lucas JW. Summary health statistics for U.S. adults: National Health Interview Survey, 2009. *Vital Health Stat 10.* 2010;(249):1–207.
4. Healthdata. Smoking in the Kingdom of Saudi Arabia: Findings from the Saudi Health Interview Survey. Access 13 November; 2021. Available: <http://www.healthdata.org/sites/default/files/files/Projects/KSA/Smoking-KSA-Findings-from-the-Saudi-Health-Interview-Survey.pdf>.
5. United States Public Health Service Office of the Surgeon General, & National Center for Chronic Disease Prevention and Health Promotion (US) Office on Smoking and Health. Smoking Cessation: A Report of the Surgeon General. US Department of Health and Human Services; 2020.
6. Babb S, Malarcher A, Schauer G, Asman K, Jamal A. Quitting Smoking Among Adults—United States, 2000–2015. *Morb Mortal Wkly Rep.* 2017;65(52):1457-64.
7. Creamer MR, Wang TW, Babb S, Cullen KA, Day H, Willis G, et al. Tobacco Product Use and Cessation Indicators Among Adults — United States, 2018. *MMWR Morb Mortal Wkly Rep.* 2019;68:1013–1019.
8. Verywellhealth. What Is Nicotine?. Access 13 November 2021. Available: <https://www.verywellhealth.com/what-is-nicotine-5075412>.
9. Drugabuse.gov. Is nicotine addictive?. Access 13 November 2021. Available: <https://www.drugabuse.gov/publications/research-reports/tobacco-nicotine-e-cigarettes/nicotine-addictive>.
10. Verywellmind. What to Know About Nicotine Use. Access 13 November 2021. Available: <https://www.verywellmind.com/nicotine-addiction-101-2825018>.
11. Medlineplus. Varenicline. Access 13 November; 2021. Available: <https://medlineplus.gov/druginfo/meds/a606024.html>.
12. Drugbank. Varenicline. Access 13 November; 2021. Available: <https://go.drugbank.com/drugs/D01273>.
13. Webmd. Varenicline 1 Mg Tablet Smoking Deterrents - Nicotinic Or Bupropion (NDRI)-Type - Uses, Side Effects, and More. Access 13 November 2021. Available: <https://www.webmd.com/drugs/2/drug-144328/varenicline-oral/details>.
14. Medscape. Varenicline. Access 13 November 2021. Available: <https://reference.medscape.com/drug/chantix-varenicline-343212>.
15. Fiore MC, Bailey WC, Cohen SJ, Dorfman SF, Goldstein MG, Gritz ER, et al. Treating Tobacco Use and Dependence: Clinical Practice Guideline. Rockville, MD: US Department of Health and Human Services, Public Health Service; 2000.
16. Steinberg MB, Akincigil A, Delnevo CD, Crystal S, Carson JL. Gender and age disparities for smoking cessation treatment. *Am J Prevent. Med.* 2006;30(5): 405-12.
17. Prochaska JJ, Benowitz NL. The past, present, and future of nicotine addiction therapy. *Annu. Rev. Med.* 2016;67:467–86.
18. Caraballo RS, Shafer PR, Patel D, Davis KC, McAfee TA. Quit methods used by U.S. adult cigarette smokers, 2014–2016. *Prev. Chronic Dis.* 2017;14:160600.
19. Jamal A, Phillips E, Gentzke AS, Homa DM, Babb SD, King BA, et al. Current cigarette smoking among adults— United States, 2016. *Morb Mortal Wkly Rep.* 2018;67(2):53–9.
20. DiGiulio A, Jump Z, Yu A, Babb S, Schecter A, Williams KS, et al. State Medicaid coverage for tobacco cessation treatments and barriers to accessing treatments—United States, 2015–2017. *Morb Mortal Wkly Rep.* 2018;67(13): 390–5.

21. Fiore MC, Jaén CR, Baker TB, Bailey WC, Benowitz NL, Curry SJ, et al. Treating Tobacco Use and Dependence: 2008 Update. U.S. Public Health Service Clinical Practice Guideline. Rockville (MD): U.S. Department of Health and Human Services, 2008.
22. U.S. Department of Health and Human Services. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. Atlanta (GA): U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014.
23. Campaign for Tobacco-Free Kids. Broken Promises to Our Children: A State-by-State Look at the 1998 Tobacco Settlement 20 Years Later, 2018. Access 13 November; 2021.
24. Ebbert J, Jimenez-Ruiz C, Dutro MP, Fisher M, Li J, Hays JT. Frequently Reported Adverse Events With Smoking Cessation Medications: Post Hoc Analysis of a Randomized Trial. *Mayo Clin. Proc.* 2021;96(7): 1801-1811.
25. Cancer.org. Prescription Medicines to Help You Quit Tobacco. Access 13 November; 2021. Available: <https://www.cancer.org/healthy/stay-away-from-tobacco/guide-quitting-smoking/prescription-drugs-to-help-you-quit-smoking.html>.
26. Rxlist. How Do Smoking Cessation Aids Work?. Access 13 November 2021. Available: https://www.rxlist.com/smoking_cessation_aids/drug-class.htm.

© 2021 Ahmed et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/77737>*