



Vulvar Edema with Severe Preeclampsia: Case Report

**Nada Douraidi ^{a*}, Sara Chater ^a, Khaoula Laaboub ^a, Aicha Kharbach ^a,
Amina Lakhdar ^a and Aziz Baidada ^a**

^a *Department of Gynecology-Obstetrics and Endocrinology, Maternity Soussi,
University Hospital Center IBN SINA, University Mohammed 5, Rabat, Morocco.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/90743>

Case Study

**Received 12 June 2022
Accepted 16 August 2022
Published 07 September 2022**

ABSTRACT

Massive vulvar edema during pregnancy is usually benign. It is often associated with patient discomfort and it is important to find the cause, as the management is influenced by the cause of edema. In this report, we describe a case of vulvar edema in a gravid preeclamptic patient. We also discussed the causes and treatment of vulvar edema in pregnancy.

Keywords: Vulvar edema; preeclampsia; external genitalia.

1. INTRODUCTION

“Several cases of the massive vulval oedema has been described in the literature associated with, diabetes, pre-eclampsia, vulvo-vaginitis, hypoproteinemia, and multiple pregnancy” [1-2]. “Very rarely is the occurrence after tocolytic therapy. Preeclampsia is a multisystemic disorder of pregnancy. It complicates 3% to 5% of pregnancies” [3]. In this report, we describe a

case of vulvar edema in a gravid preeclamptic patient in a case in the gynecological and obstetrical department the Souissi maternity hospital Rabat.

2. CASE REPORT

A 19-year-old patient, gravida (G)1, para 1, G1 current pregnancy, which was estimated at 34 weeks, and 4 days. She was transferred from a

*Corresponding author: E-mail: nadadouraidi49@gmail.com;

peripheral hospital to our tertiary care center with a main complaint of swelling of external genitalia for four days and pain. There was no other significant past, obstetric, or surgical history.

On examination, the patient was afebrile, the blood pressure was 160/90 mm Hg and pulse rate was 86 per min. Obstetric examination revealed the fetus was in cephalic presentation, the fundal height was 30 cm and normal active fetal heart rate. Pelvic examination including inspection of the vulvar showed a massive bilateral vulvar edema more marked on the right side (Fig. 1) and moderate pedal edema beyond the ankles. Obstetric ultrasound revealed a progressive monofetal pregnancy in cephalic presentation fundic placenta estimated at 2200g.

The laboratory exams revealed a positive proteinuria at 3+, The 24 h urine collection showed significant proteinuria at 4.4 g/24 H. The Other laboratory tests were within the normal range. The diagnosis of severe preeclampsia was made.

Anti-Hypertensive treatment with methyldopa and magnesium sulphate to prevent convulsion. Administration of steroid to accelerate fetal lung maturity and bed rest. After six hours of monitoring and despite all these measures and maximum dose of anti-hypertensive treatment, the patient kept a high blood pressure 180/110 mmHg and a severe headache with a nonreactive fetal heart rate pattern.

An emergency C-section was indicated resulted in the delivery of a live born female of 2140g grams and Apgar score 6-10 at first and fifth minutes, respectively. During the operation, it was aspirated almost 600 cc's of ascites. Two weeks into the postpartum monitoring, the patient showed regression of her vulvar edema with stabilization of blood pressure and negative proteinuria.

3. DISCUSSION

Vulvar edema is associated with a variety of conditions. The edema can result from inflammatory conditions, infections, infestations, trauma, pregnancy, tumors and iatrogenic causes. At times, it is difficult to determine the cause of the vulvar edema.

“This condition should be taken cautiously since it might be caused by pre-eclampsia or diabetes” [4]. “During pregnancy, the significant increase in blood volume, blood pressure, and vascular permeability, coupled with compression of the inferior vena cava by the growing uterus, can cause edema. While the exact pathophysiology of preeclampsia remains poorly understood, the disease is characterized by both systemic and localized vascular endothelial dysfunction. This is demonstrated focally in the pulmonary edema, cerebral edema, and hepatic capsule rupture and failure, all of which are significant risks of preeclampsia” [5].



Fig. 1. Vulvar edema

As for the modalities of management, a surgical treatment by draining the edema [6-7] and in some cases is management that is more conservative [8] in most cases resolves spontaneously after delivery such as our case report.

“Treatment of vulvar edema is necessary, since it is painful, uncomfortable and may cause occlusion of the vaginal outlet and makes a vaginal delivery extremely painful with a high risk of infection of the edema” [9]. “If a conservative treatment is indicated for vulvar edema, a symptomatic treatment should be instituted which include bed rest, Trendelenburg positioning, and application of ice bags, hypertonic saline bags and local antibiotics, with a close monitoring of the evolution of the edema” [10].

4. CONCLUSION

Vulvar edoema should also be taken into consideration as an important marker for severe preeclampsia as spontaneous enormous vulvar edoema is uncommon during pregnancy and poses management issues as well as significant patient discomfort, notably pain. The management of the vulvar edema depends on the cause and its clinical impact.

CONSENT

An informed consent was obtained from the patient for this publication.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Reynolds D. Severe gestational edema. *Journal of Midwifery Woman's Health*. 2003;48:146-148.
2. Saha PK. Massive vulval edema in pregnancy. *Internet Journal of Gynaecology and Obstetrics*. 2007;6.
3. Bokslag A, Van Weissenbruch M, Mol BW, De Groot CJ. Preeclampsia; short and longterm consequences for mother and neonate. *Early Human Development*. 2016;102:47-50. PubMed Google Scholar
4. Bush JP. Oedema of vulva due to toxemia of pregnancy. *British Medical Journal*. 1946;2:988.
5. Houben AJHM, De Leeuw PW, Peeters LLH. Configuration of the microcirculation in pre-eclampsia: possible role of the venular system. *Journal of Hypertension*. 2007;25(8):1665–1670.
6. Radomański T, Sikorski R, Baszak E. Vulvar edema in pregnancy. *Ginekologia Polska*. 1998;69(12):1067–1070.
7. Lindsey JS, De Vente JE. Surgical management of massive labial edema in a gravid preeclamptic diabetic. *Case Reports in Obstetrics and Gynecology*; 2014.
8. Deren O, Bildirici I, Al A. Massive vulvar edema complicating a diabetic pregnancy. *European Journal of Obstetrics and Gynecology and Reproductive Biology*. 2000;93(2):209–211.
9. Trice L, Bennert H, Stubblefield PG. Massive vulvar edema complicating tocolysis in a patient with twins: a case report. *J Reprod Med*. 1996;41:121–124.
10. Guven ES, Guven S, Durukan T, Onderoglu L. Massive vulval oedema complicating pregnancy. *J Obst Gynaecol*. 2005;25:216–218.

© 2022 Douraidi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/90743>