



Integrated Nutrient Management: A Pathway to Enhanced Growth and Yield in *Amaranthus tricolor* L.

Saurabh Yadav ^{a*}, Rameshwar Jangu ^a, Suraj Malviya ^b,
Jaswan Prajapati ^a, Kalp Das ^a and K.P. Asati ^c

^a Department of Vegetable Science, College of Horticulture and Forestry, Punjab Agricultural University, Ludhiana, Punjab, 141004, India.

^b IIT, Indore, Madhya Pradesh 452001, India.

^c Department of Horticulture, College of Agriculture, Indore, Madhya Pradesh, 452001, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i71048>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118109>

Short Research Article

Received: 20/04/2024

Accepted: 21/06/2024

Published: 24/06/2024

ABSTRACT

The present study was conducted to determine the effect of different levels of manures, fertilizer, and biofertilizers application, either alone or in combination on seed germination, vigor, and yield of *Amaranthus*. Recommended rates of organic manures viz., farmyard manure (5 t/ha) and vermicompost (2.5 t/ha) biofertilizers i.e., *Azotobacter* and *Phosphate solubilizing bacteria* @ 200g per 10 kg of seed were integrated with the recommended dose of fertilizer @ 60: 40: 20 kg/ha NPK. The experiment was designed with twelve treatments with three replications that were laid out in a randomized complete block design. Results revealed that the treatment T₁₂ which consists of half dose of NPK + half dose of vermicompost + *Azotobacter* and PSB observed early emergence of

*Corresponding author: E-mail: bittu1441994@gmail.com;

Cite as: Yadav, Saurabh, Rameshwar Jangu, Suraj Malviya, Jaswan Prajapati, Kalp Das, and K.P. Asati. 2024. "Integrated Nutrient Management: A Pathway to Enhanced Growth and Yield in *Amaranthus Tricolor* L". *Journal of Advances in Biology & Biotechnology* 27 (7):885-92. <https://doi.org/10.9734/jabb/2024/v27i71048>.

seeds (3.60 days) along with higher germination percentages (84.53 percent) as well as vigorous seedlings in terms of vigor index-1 (5.75) and vigor index-2 (5.56). The crop was harvested three times at a height of 20 cm and by the application of treatment T₁₂, plants take minimum time for harvest i.e., 21.65, 30.65, and 41.16 days along with the highest yield i.e., 0.50, 1.15, and 0.85 kg per plot at first second and third harvest respectively. Therefore, it is clear that combining inorganic fertilizers with organic manures could increase soil health in addition to enhancing growth and facilitating consistent nutrition for the Amaranthus crop.

Keywords: *Amaranthus*; germination; integrated nutrient management; vigour; yield.

1. INTRODUCTION

“Vegetables are essential components of our daily diet, with a recommended intake of at least 300 grams per person per day, including 125 grams of leafy vegetables. Leafy vegetables, often dubbed 'mines of minerals,' contribute significantly to our nutritional intake, offering vitamins, minerals, protein, carotene (vitamin A), iron, and ascorbic acid (vitamin C)” [1]. “Among these, Amaranthus stands out as a widely consumed vegetable across India” [2]. Derived from the Greek word 'Anthos' meaning everlasting or unwilting, Amaranthus, also known as 'Chaulai' or the 'poor man's vegetable,' belongs to the genus Amaranthus, family Amaranthaceae, which comprises 65 genera and 850 species worldwide. *Amaranthus tricolor* L., extensively cultivated in Bengal historically [3], holds cultural and agricultural significance in South India as well [4,5]. Originating in India [6], Amaranthus demonstrates centers of diversity in Central and South America, India, and Southeast Asia, with secondary diversity in West Africa and East Africa [7]. Its attributes as a C4 plant, coupled with its short-duration growth and high yield per unit area, make it suitable for crop rotation. The nutritive value of Amaranthus is noteworthy, offering energy, carbohydrates, fat, protein, vitamins, and trace elements [8-11]. However, its performance hinges directly on soil nutrient status managed through organic and inorganic fertilizers. While chemical fertilizers have been traditionally employed, their adverse effects on soil health, ecosystem balance, and human health, coupled with escalating costs and availability issues, necessitate the exploration of alternative nutrient sources. Organic manures, composts, vermicomposts, crop residues, green manures, and biofertilizers present viable alternatives, offering cost-effectiveness and availability. Integrated Nutrient Management (INM) emerges as a promising approach, leveraging organic and chemical fertilizers alongside biofertilizers to sustain crop yield while enhancing fertilizer efficiency and soil health [12].

This study, titled “Integrated Nutrient Management: A Pathway to Enhanced Growth and Yield in *Amaranthus tricolor* L.,” aims to explore this approach, recognizing the diversity in nutrient sources and their implications for improved crop performance.

2. MATERIALS AND METHODS

The experiment was conducted during the *kharif* season at the Nursery area, Department of Horticulture, College of Agriculture, Indore, Madhya Pradesh, India. The experiment field was prepared by two deep ploughing with disc plough and planking to bring the field to a good tilth condition. Each experimental unit was demarcated and the layout was drawn as per the plan finally, the field was divided into 39 plots (2.25 m X 1.05 m each plot) having irrigation channels and paths. There were thirteen treatments and each treatment was allocated randomly in each plot. The thirteen treatments, T1: Control (No use of Fertilizers and manures), T2: Recommended dose of NPK (60,40,20), T3: 100% FYM (5 tonnes/ha), T4: 100% Vermicompost (2.5 tonnes/ha), T5: Azotobacter (200g/10 kg seed) + PSB (200g/10 kg seed), T6: 75% Recommended dose of fertilizers + 25% Vermicompost, T7: 75% Recommended dose of fertilizers + 25% FYM, T8: 75% Recommended dose of fertilizers + 25% Vermicompost + Azotobacter + PSB, T9: 75% Recommended dose of fertilizers + 25% FYM + Azotobacter + PSB, T10: 50% Recommended dose of fertilizers + 50% Vermicompost, T11: 50% Recommended dose of fertilizers + 50% FYM, T12: 50% Recommended dose of fertilizers + 50% Vermicompost + Azotobacter + PSB, T13: 50% Recommended dose of fertilizers + 50% FYM + Azotobacter + PSB were evaluated in randomized block design with the three replications. “Before sowing in the plots, seeds were treated with thiram @ 2.5 g/kg and then with Azotobacter and Phosphate solubilizing bacteria @ 200g/10 kg seed. FYM @ 5 tonnes/ha and vermicompost @ 2.5 tonnes/ha were

incorporated in the field at the time of field preparation as per treatments. Half dose of nitrogen @ 60 kg/ha and a full dose of phosphorus @ 40 kg/ha and potassium @ 20 kg/ha was also given as per treatment in the plots as basal dose and the remaining quantity of nitrogen was applied as top dressing at each foliage cutting. By maintaining row to row distance of 45 cm, seeds were sown in the plots @ 2 kg/ha in the third week of July. Gradually thinning was done at 5 cm height to achieve a final spacing of 15 cm between the plants. All cultural operations were performed as per recommendations. The statistical analysis was carried out as per the procedure” [13].

For measuring Days to the first seedling emergence daily observations were taken, until the first foliage leaf appeared. Germination percent was recorded according to ISTA, 2011. Vigour index I and II were computed using the formula [14], and expressed in the whole number. Five random plants were selected and properly tagged in each replication for every treatment to collect the data related to the number of days taken to achieve 20 cm height as harvesting of greens were done at 20 cm height. The greens obtained from each plot and replication at the first, second, and third harvest at 20 cm height were weighted and the Yield /plot (kg) was worked out.

Germination percentage

$$= \frac{\text{Number of germinating seeds} \times 100}{\text{Total number of seeds}}$$

Vigour Index – 1

$$= \frac{\text{Germination percentage} \times \text{Seedling length (cm)}}{100}$$

Vigour Index – 2

$$= \text{Seedling dry weight (g)} \times \text{Germination percentage}$$

3. RESULTS

The results of the experiments are presented in Table 1. There is a significant effect of Integrated Nutrient Management on all the traits viz. Number of days taken for germination, germination percentage, Vigor index-1 and-2, number of days taken to harvest, and yield (kg per plot). By the application of treatment combination T₁₂ seeds took of 3.60 days for germination, which was followed by 4.10 days due to the treatment combination T₁₃ while a maximum of 6.51 days was recorded due to the application treatment T₁ (Fig. 2). Similarly, a maximum 84.53 germination percentage was also observed due to the application of treatment T₁₂ which was succeeded by 78.19 percent by the use of treatment T₁₃ meanwhile the minimum of 58.74 percent germination was recorded in treatment T₁ (Fig. 1). Following this, the maximum value of vigor index-1 (5.75) and vigor index-2 (5.56) was also observed in T₁₂ followed by the 5.38 (vigor index-1) and 4.85 (vigor index-2) by the application of T₁₃ (Fig. 1). At the same time, minimum values of vigor index-1 (1.93) and Vigor index- (1.01) were observed by the use of treatment T₁ (control) (Fig. 1). Moreover, by the use of treatment T₁₂, plant took a minimum of 21.65, 30.65, and 41.16 days to attain 20 cm height for all three consecutive first, second, and third harvest respectively (Fig. 2). It was followed by the application of treatment T₁₃ by which plants took 23.69, 32.9 and 43.86 days for first, second and third harvest respectively at 20 cm height (Fig. 2). While the application of treatment




Fig. 1. Effect of different treatment combinations on Vigour index-, Vigour Index-2 and Germination percentage




Fig. 2. Effect of different treatment combinations on number of days taken to germination along with Days taken to first, second, and third harvest




Fig. 3. Effect of different treatment combinations on yield (kg/plot) at 20 cm height at first, second, and third harvest

T₁ took maximum 32.25, 44.90 and 59.84 days for all the three harvests respectively (Fig. 2). Similarly, a maximum yield of 0.50 kg per plot, 1.15 kg per plot, and 0.85 kg per plot was also observed by the application of treatment T₁₂ at all three harvests (Fig. 3). This was succeeded by T₁₃ by which 0.38 kg per plot, 0.99 kg per plot, and 0.74 kg per plot yield was obtained at each harvest (Fig. 3) while the application of treatment T₁ yielded a minimum of 0.15 kg per plot, 0.20 kg per plot, and 0.18 kg per plot greens at each harvest respectively (Fig. 3).

4. DISCUSSION

The overall performance of Amaranthus is better in the treatment combination T₁₂. It might be due

to the positive response of the crop toward the nutrition available from different sources like fertilizer, vermicompost, and biofertilizers. The availability of effective microorganisms through biofertilizers along with nutrients supplied by vermicompost and chemical fertilizers which change the soil plant ecosystem in terms of soil health and growth and development of the plant consequently increased the yield and its attributing traits of Amaranthus [15,16]. Biofertilizers stimulate plant growth by providing necessary nutrients as a result of their colonization at their rhizosphere (*Azotobacter*, *Azospirillum*, *Pseudomonas*, *Phosphate solubilizing bacteria*, and *Cyanobacteria*) or by symbiotic association (*Rhizobium*, *Mycorrhizae* or *Frankia*) [17].

Table 1. Effect of integrated nutrient management on days taken for germination, Germination percentage, Vigour index – 1, Vigour index – 2, number of days taken to harvest, yield kg/plot

Treatments	Days taken for Germination	Germination Percentage	Vigour Index - 1	Vigour Index - 2	Number of Days taken at 20 cm Height			Yield kg/plot at 20 cm Height		
					1 st harvest	2 nd harvest	3 rd harvest	1 st harvest	2 nd harvest	3 rd harvest
T ₁	6.51	58.74	1.93	1.01	32.25	44.90	59.84	0.15	0.20	0.18
T ₂	6.10	63.48	2.95	2.24	28.11	40.01	55.64	0.22	0.58	0.40
T ₃	4.82	66.20	2.87	2.58	29.75	41.11	52.77	0.21	0.55	0.42
T ₄	4.50	70.12	3.49	3.25	28.51	40.68	50.89	0.24	0.61	0.48
T ₅	4.79	68.24	3.46	2.79	29.12	41.7	52.59	0.22	0.58	0.41
T ₆	5.81	72.91	4.03	3.65	27.68	38.79	50.09	0.27	0.72	0.49
T ₇	5.90	69.63	3.61	2.97	27.84	38.97	49.89	0.25	0.63	0.47
T ₈	4.33	77.35	4.94	4.51	26.20	36.38	48.26	0.34	0.87	0.62
T ₉	4.57	75.61	4.58	4.12	26.86	35.85	48.58	0.31	0.83	0.70
T ₁₀	5.01	72.02	4.39	3.74	26.56	38.51	49.01	0.30	0.79	0.53
T ₁₁	5.36	73.85	4.23	3.69	27.77	39.14	49.61	0.28	0.77	0.51
T ₁₂	3.60	84.53	5.75	5.56	21.65	30.65	41.16	0.50	1.15	0.85
T ₁₃	4.10	78.19	5.38	4.85	23.69	32.9	43.86	0.38	0.99	0.74
S. Em. ±	0.16	2.11	0.12	0.20	0.81	1.13	1.46	0.01	0.02	0.01
CD at 5%	0.47	6.17	0.36	0.59	2.37	3.29	4.26	0.02	0.06	0.04

Treatment T₁₂ has many growth-promoting hormones like indole -3- acetic acid (IAA), cytokinin, gibberellins, etc., and humic acid which could have been responsible for the faster germination of seed. Biofertilizers also promote seed germination and early development by excreting phytohormones such as auxins and gibberellins, etc. Besides, during metabolism the bacteria excrete organic acids (citric acid, malic acid, etc.) as well, thus helping nutrient uptake at a later stage of growth. It was reported that 10g inoculation of *Azotobacter* showed 100 percent germination and also reduced the average time taken to start germination [18]. This result is in agreement with the previous findings [19,20] in *Amaranthus*. Additionally, Nitrogen especially nitrate stimulates the germination of *Amaranthus* [21,22]. Previously it was reported that many weed species including *Amaranthus* nitrogenous compounds stimulate germination [23]. Similar findings in variations in germination percentage were also confirmed by previous researchers in *Amaranthus* [24,20].

Vermicompost along with Biofertilizers produce or change the concentration of plant hormones, asymbiotic nitrogen fixation, and solubilization of mineral phosphate and other nutrients during germination which improves the seed vigor. This was in line with the previous findings [25-27] which stated that the seedling vigor index -1 and vigor index -2 increased with the combined application of inorganic fertilizers, organic manures, and biofertilizers.

Moreover, Biofertilizers and organic manures enrich the soil with organic matter content which increases the porosity of the soil which is beneficial to provide a congenial environment to roots for uptake of the available nutrients from the soil. The adequacy of manure decreased the number of days from planting to the first harvest and increased the number of harvests before senescence [28]. These findings are in accordance with previous researchers who observed a higher dose of nitrogen percentage of 20 % N from castor cake + 20 % N from vermicompost + 30 % N from poultry manure + 30 % N from FYM early harvest of the crop was possible [29].

Enhanced nitrogen content encourages increased growth and yield of crops [30,31]. Since it is the main yield factor and is considered the characteristic constituent of functional plasma, an integral part of chlorophyll molecules, proteins, amino acids nucleic acids (RNA and

DNA), nucleotides, phosphatides, alkaloids, enzymes, coenzymes, hormones, vitamins [32]. Due to the application of nitrogen through organic manures, inorganic fertilizers, and biofertilizers, plants received a sufficient amount of other essential elements throughout their growth period. They nourished properly which improved the green leaves yield in the first cutting as well as in subsequent cuttings (second and third). Interestingly it is clear from Fig. 3, that the yield at the second harvest is maximum in all the treatment combinations, It might be because they encouraged the vigorous growth of the root system which ultimately helped in better absorption and utilization of nutrients from soil solution as a result of high dry matter accumulation took place and their translocation in plant parts, which in turn reflected in better plant growth [29]. These results are similar to the previous findings in *Amaranthus* [19].

5. CONCLUSION

Based on the aforementioned findings, it can be said that application of N: P: K @ 30:20:10 + half dose of vermicompost @1.25 tonnes per ha + *Azotobacter* @ 200 g/10kg seeds and PSB @ 200 g/10kg seeds was significantly effective in improving the germination, vigor, and yield of *Amaranthus* along with soil physical conditions.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Van Soest PJ, Robertson JB, Lewis BA. Symposium: carbohydrate methodology, metabolism and nutritional implications in dairy cattle. *Journal of Dairy Sciences*. 1991;74:3583-97.
2. Akubugwo IE, Obasi NA, Chionyere GC, Vgbogo AE. Nutritional and chemical value of *Amaranthus hybridus* L. leaves from Afikpo, Nigeria. *African Journal of Biotechnology*. 2007;6(24):2833-2839.

3. Watts G. A Dictionary of the Economic Products of India, Gordhan and Company, Delhi. 1989;1:204-216.
4. Sankaran A. Cultivation of leafy vegetables in the northern circars. Madras Agricultural Journal. 1943;31:285-286.
5. Choudhary R, Diwakar AK, Meena AK, Shukla S, Prakash C. Assessment of fertility status of soils under different cropping patterns in rainfed semi-arid eastern plain of Rajasthan, India. International Journal of Environment and Climate Change. 2024;14(5):192-204.
6. Nath P. Vegetables for the Tropical Region, ICAR, New Delhi. In: C.R. Muthukrishnan and I. Irulappan. Amaranthus. T.K. Bose and M.G. Som (eds.). Vegetable crops in India. Naya Prokash, Calcutta six, 1976;670.
7. Grubben GJH. Tropical vegetables and their genetic resources, (H.D. Tindall and J.T. Williams eds.) Rome. 1977;91-110.
8. Kumar CR, Jacob J, Kiran. Effect of different organic media and nutrient management on growth and yield of Amaranthus on terrace gardens. Trends in Biosciences. 2015;8(9):2433-37.
9. Kaur G, Singh I, Behl R, Dhankar A. Effect of Different Integrated Nutrient Management Approaches on Growth, Yield Attributes and Yield of Wheat (*Triticum aestivum* L.) Crop: A Review . Asian Journal of Soil Science and Plant Nutrition. 2024;10(1):457-468.
Available:<https://doi.org/10.9734/ajsspn/2024/v10i1252>
10. Vishwakarma M, Chauhan MVR. The impact of integrated nutrient management on growth and yield of palak (Indian Spinach). International Journal of Plant & Soil Science. 2023;35(20):763-772.
Available:<https://doi.org/10.9734/ijpss/2023/v35i203863>
11. Gopaldasundaram P, Bhaskaran A, Rakkiyappan P. Integrated nutrient management in sugarcane. Sugar Tech. 2012;14:3-20.
12. Singh GB, Biswas PP. Balanced and integrated nutrient management for sustainable crop production. Indian Journal of Fertilisers. 2000;45(5):55-60
13. Panse VG, Sukhatme PV. Statistical Methods for Agricultural Workers. Indian Council for Agricultural Research, New Delhi. 1989;359
14. Abdul-Baki AA, Anderson JD. Relationship between decarboxylation of glutamic acid and vigor in soybean seed 1. Crop science. 1973;13(2):227-32.
15. Yadav S, Asati KP, Barche S. Effect of integrated nutrient management strategies on growth and yield of Amaranthus (*Amaranthus tricolor* L.). The Pharma Innovation Journal. 2022;11:2545-9.
16. Choudhary R, Nath T, Sharma PK, Nagar PK, Meena AK, Naga IR. Soil quality assessment of different villages of malpura block in Tonk District of Rajasthan, India. International Journal of Plant & Soil Science. 2024;36(5):352-362.
17. Varma A, Shuepp H. Mycorrhization of the commercially important micro propagated plants. Critical Reviews in Biotechnology. 1995;15:313-28.
18. Monika K, Dheeraj V. Effect of biofertilizers (*Azotobacter* and *PSB*) and their combinations on germination and survival of *Lens culinaris* Medic. Environment and Ecology. 2010;28(1B):703-5.
19. Tayade MS, Bawkar SO, Deshmukh UB. Integrated nutrient management in Amaranthus (*Amaranthus tricolor* L.). Asian Journal of Horticulture. 2012;7(2):291-93.
20. Shanker Y. Effect of *Azadirachta indica* vermicompost on growth of Amaranthus and Fenugreek. SIRJ-AZASN 2014; 1(1): 34-39.
21. Blackshaw RE, Rode LM. Effect of ensiling and rumen digestion by cattle on weed seed viability. Weed Science. 1991;39(1): 104-8.
22. Hernández JA, Díaz-Vivancos P, Acosta-Motos JR, BarbaEspin G. Potassium nitrate treatment is associated with modulation of seed water uptake, antioxidative metabolism and Phytohormone levels of Pea Seedlings. Seeds. 2021;1, 5-15.
23. Baskin JM, Baskin CC. Role of temperature in the germination ecology of three summer annual weeds. Oecologia. 1977;30:377-82.
24. Charachimwe R, Tembo L, Mhungu S. Response of pig weed Amaranthus cruentus to organic and inorganic fertilizers. International Journal of Agronomy and Agricultural Research. 2018;13(1):1-10.
25. Wani IM, Narayan S, Magray MM, Khan F. A, Mir SA, Yaseen I, Khurshid A. Effect of organic manures and biofertilizers on seed quality of radish (*Raphanus sativus* L.)

- under Temperate Conditions of Kashmir. The Pharma Innovation Journal. 2023;15:315-319.
26. Maheshbabu HM, Ravi Hunje RH, Patil NB, Babalad HB. Effect of organic manures on plant growth, seed yield and quality of soybean. Karnataka Journal Agriculture Sciences. 2008;21(2):219-21.
27. Renugadevi J, Balamurugan P. Effect of bio-fertilizer in seed yield and seed quality in sunflower. Madras Agricultural Journal. 2002;89(1/3):135-7.
28. Ojo OD, Olufolaji AO. The Effects of different phosphorus sources on growth and yield of grain amaranth (*Amaranthus cruentus* L.). In Proceedings of the 15 Conference of the Horticultural Society of Nigeria. 1997;53-55.
29. Acharya MD, Patel GS, More SG, Chaudhary BM, Rabari SS. Effect of different organic sources of nitrogen on growth parameters and yield attributes of amaranth (*Amaranthus* Spp.) Cv. Arka Suguna. International Journal of Chemical Studies. 2008;6(3): 575-579.
30. Jangu R, Meena ML, Shivran BC, Chand R. Determining the combined effect of nitrogen and spacing on growth parameters in lettuce (*Lactuca sativa* L.). The Pharma Innovation Journal. 2022; 11(2):1740-45
31. Wu K, Wang S, Song W, Zhang J, Wang Y, Liu Q, Fu X. Enhanced sustainable green revolution yield via nitrogen-responsive chromatin modulation in rice. Science. 2020;367(6478):eaaz2046.
32. Castellanos J, Uvalle-Bueno J, Aguilar-Santielises A. Manual de interpretación de análisis de suelos y aguas. 2a Edición. San Miguel de Allende, Guanajuato, México: Instituto de Capacitación para la Productividad Agrícola (INCAPA). 2000; 226.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/118109>