



# Recycling Selected Wetlands' Sediments for a Vegetable Crop Cultivation in Terrestrial Ecosystem of a Humid Region

Md. Shahadat Hossain <sup>a\*</sup>, Md. Abul Kashem <sup>a</sup>, Israt Jahan <sup>a</sup>  
and Kazi Sanjida Begum <sup>a</sup>

<sup>a</sup> Department of Soil Science, Sylhet Agricultural University, Bangladesh.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ajsspn/2024/v10i3344>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120669>

Original Research Article

Received: 22/05/2024

Accepted: 25/07/2024

Published: 31/07/2024

## ABSTRACT

This study aimed to evaluate the selected wetlands' sediments regarding physicochemical attributes including heavy metals (HMs), and their potentiality to be used in crop cultivation with or without mixing with different generic farmers' field soils in the upland. Forty geo-referenced sediment samples were collected from each study site (n=3), and mixed to get three separate composite samples for mineral nutrients, HMs analyses and experimentation. Two pot experiments were conducted following Completely Randomized Design (CRD) with three replicates using farmers' field soils (FFS) Gleysols and Cambisols to grow amaranth. The treatments were: T<sub>0</sub>: FFS; T<sub>1</sub>: FFS + wetland sediment of hilly area I (1:1); T<sub>2</sub>: FFS + wetland sediment of hilly area II (1:1); T<sub>3</sub>: FFS + wetland sediment of a floodplain area (1:1); T<sub>4</sub>: wetland sediment of hilly area I; T<sub>5</sub>: wetland

\*Corresponding author: Email: [mshossain.ss@sau.ac.bd](mailto:mshossain.ss@sau.ac.bd);

**Cite as:** Hossain, Md. Shahadat, Md. Abul Kashem, Israt Jahan, and Kazi Sanjida Begum. 2024. "Recycling Selected Wetlands' Sediments for a Vegetable Crop Cultivation in Terrestrial Ecosystem of a Humid Region". *Asian Journal of Soil Science and Plant Nutrition* 10 (3):327-42. <https://doi.org/10.9734/ajsspn/2024/v10i3344>.

sediment of hilly area II; T<sub>6</sub>: wetland sediment of a floodplain area. Regarding nutrient elements, exchangeable K, extractable P, Ca, and Zn were observed higher in wetland sediment of hilly area II except organic carbon (OC). Only Cd, Pb, and Ni were individually found higher than the maximum permissible addition for all wetland sediments, but all HMs collectively posed no “ecological risk”. The highest values of the plant growth index of amaranth were observed for T<sub>5</sub> treatment. Moreover, the highest total fresh and dry biomasses were found in T<sub>5</sub> treatment. The highest nutrient uptake was observed in T<sub>5</sub> treatment for the most of nutrients. Studied wetlands’ soil materials, OC, and nutrients could be used to benefit upland crop cultivation.

**Keywords:** Sediment; heavy metals; gleysols; cambisols; nutrient uptake.

## 1. INTRODUCTION

Wetlands are the lands that remain under water either permanently for years or decades, or temporarily in seasons of a year. A wetland possesses a distinct ecosystem that serves as a home for diverse plant and animal species [1]. Wetlands vary greatly due to local and regional differences in natural (soil, topography, climate, hydrology, water chemistry, vegetation etc.) and anthropogenic (human interference) factors, and they are broadly categorized as ‘tidal’ and ‘non-tidal (inland)’ wetlands [2]. Water in wetlands is being threatened quantitatively and qualitatively by the sedimentation of particulate matters including soil materials. As a result, wetland’s retention capacity can be lowered down [3], reservoirs’ bed can be raised up [3], evaporation losses can be increased [4], and eutrophication processes can be accelerated [5]. So, it is crucial to reclaim the wetlands to protect the water supply and biodiversity through excavating the bottom bed materials. Dredging is an operative method for excavating and re-excavating land to improve the wetland’s water bearing ability [6,7]. Globally, millions of tons of dredged sediment are produced by incessant dredging. For instance, around 200 m<sup>3</sup> of dredged material are thought to be produced annually in Europe [8, 9, 10].

Wetland’s bottom sediment (soil) is an indispensable part of an ecosystem [11] and possesses distinct hydrology and unique physical setting. Its physicochemical and biological characteristics are vital anthropopressure indicator [12,13]. Sediments can improve terrestrial soil pH [12,14,15,16], soil aggregate [3,17], soil organic carbon [17,18], soil total N, extractable P [19], Exchangeable K, microbial biomass [20], sorption properties [17,21] and crop productivity [21,22]. Considering the larger issue of global food security and soil sustainability, wetlands’ sediment availability as

fertilizers can be a key component in upland agriculture.

Many researchers [14,17,23,24,25] underscored the recycled use of excavated sediment for terrestrial agriculture. This usage of sediment is recommended only when the toxic contaminant content is the below maximum permissible limit [26,27,28]. Before utilization, their evaluations in terms of the nature and quality are prerequisites [29,30], because wetland sediment is influenced by numerous variables. These include (i) the nature and quality of stream water feeding wetland (ii) anthropogenic and agricultural activities (fertilizer, pesticide etc. applications) in the wetland and adjoining terrestrial areas (iii) the nature of adjacent landscape (hills or plain land) and soils (alluvial or colluvial) (iv) Runoff water quality and quantity reaching the water body, and (v) Openness of wetland, etc. Therefore, in the light of above discussion it can be said that inland wetlands as well as their sediments vary from one to another. Furthermore, the responsible factors can give a unique make up of sediment in certain wetlands in a region. In addition, in the areas like our study sites where flood water visits in the wet season, previously collected sediment could be used for pot culturing of vegetables to meet up the daily needs of affected people. In humid sub-tropics, runoff induced uni-directional loss-gain phenomenon of soil materials between laterally embedded terrestrial land and wetland can be transformed into reversible phenomenon (a sustainable approach) through reuse of the wetland’s bottom sediment in terrestrial upland. The evaluation of the diverse sediments and their utilization in different generic terrestrial upland might give encouraging and/or different results. Therefore, the present study has been designed to evaluate the selected wetlands’ sediments regarding physicochemical attributes including heavy metals, and their potentiality to be used in stem amaranth cultivation with or without mixing

with different generic farmers' field soils in terrestrial ecosystem.

## 2. MATERIALS AND METHODS

Two studies were performed to address the objectives of this research work.

**Study 1:** Assessment of physicochemical attributes and heavy metal contaminants in three unique wetlands' sediment

**Study areas:** Three unique and genetically different wetlands (two haos and a beel) of Bangladesh were the study areas of this research work (Fig. 1). One of the study sites is the wetland of hilly area I (locally known as Hakaluki haor) which is the largest wetland in Bangladesh, located in the adjacent to Assam border close to hills. The rapid degradation of the ecology of this land is causing devastating consequences on the community surrounding the land. Its home country Bangladesh has declared it as an "Ecologically Critical Area" in 1999 [31]. Another site is wetland of hilly area II (locally known as Tanguar Haor) which is a unique wetland ecosystem located in Bangladesh, adjacent to Indian hills which bears natural importance and serves as the source of livelihood for more than 40,000 people. The host country Bangladesh has also declared it as an "Ecologically Critical Area" in 1999 for its overexploitation (www.wikipedia.org.). In addition to foreign hills (India, Assam), the adjoining area in Bangladesh (for wetland of hilly areas I and II) is characterized by the presence of numerous small hills (locally called "tillas"). The third site is wetland of floodplain area (locally known as Chalan Beel) which is one of the largest and richest non-tidal wetlands of Bangladesh. It comprises a series of depression interconnected by various channels. This marshy land is being silted up rapidly [32].

**Soil samples collection, processing and analysis:** Forty geo-referenced soil samples (sediment) with the assistance of a GPS device (0-20 cm, around 3 kg each) from each site were collected in a zig-zag random fashion maintaining a distance of around half a kilometer between sampling points (Fig. 1). Composite soil samples were obtained by mixing collected samples for each site. Sub-sampling was done from each composite sample, and they were processed for further analysis (K, P, S, Ca, Mg, Zn, Fe, Mn, Cu, Cd, Pb and Ni) by using standard methodology (SI 1). Remaining soils were used as a media for pot culture.

**Non-specific biomass targeted soil/sediment pollution risk by heavy metals (Zc):** Total heavy metal pollution index (Zc) was counted by the following formula [33].

$$Z_c = \sum K_{ki} - (n - 1) \dots\dots\dots (1)$$

Where  $K_{ki}$  was the ratio (from observed analytical value and reference value) of the  $i$ -th element;  $n$  was the number of elements as pollutants.

**Plant specific targeted soil/sediment pollution risk by heavy metals as nutrient element:** Total Heavy metal (nutrient) pollution index ( $Z_{c_n}$ ) was defined and found by the following formula:

$$Z_c = \{ \text{Metal (as nutrient) conc. in soil} - (\text{Critical nutrient concentration} \times 3) \} \dots\dots\dots (2)$$

**Study 02:** Assessment of sediments as media for pot culturing of a fast-growing crop stem amaranth in Eutric Gleysols and Dystic Cambisols

**Study site:** Two pot experiments were conducted at the experimental net house of the Department of Soil Science at Sylhet Agricultural University, Sylhet (Elevation: 22.56 meters, North latitude: 24°54'39.16" and East longitude: 91°54'04.79"). Sylhet region experiences average rainfall of approximately 500 mm and average temperature of 25°C during the month of August to November due to its humid subtropical monsoon environment.

**Pot cultured experiments:** Each pot (diameter 13 cm and height 16 cm) was filled with 3.3 kg of collected soil and/or sediment as stated in study 1. The experiment encompasses seven different treatments, namely:  $T_0$ : Farmer's field soil (FFS) [Eutric Gleysols for 1st pot experiment and Dystic Cambisols for 2nd pot experiment];  $T_1$ : FFS + Wetland sediment of hilly area I (1:1);  $T_2$ : FFS + Wetland sediment of hilly area II (1:1);  $T_3$ : FFS + Wetland sediment of floodplain area (1:1);  $T_4$ : Wetland sediment of hilly area I;  $T_5$ : Wetland sediment of hilly area II;  $T_6$ : Wetland sediment of floodplain area. The initial statuses of Eutric Gleysols & Dystic Cambisols are presented in Table 1.

Completely Randomized Design (CRD) was followed with three replications maintaining the space 30 cm and 30 cm, respectively, between two pots and two rows for each experiment. All experimental pots for both experiments received the recommended dose of fertilizers (Urea 195.6 and 195.6 kg ha<sup>-1</sup>, TSP 140 and 120 kg ha<sup>-1</sup>, MoP 85.254 and 90.15 kg ha<sup>-1</sup>, Gypsum 75.136

and 70.15 kg ha<sup>-1</sup> for the first and second studies, respectively) based on soil test value which was calculated by the formula from Fertilizer Recommendation Guide [34]. Based on the pot soil pH value, 1 g kg<sup>-1</sup> soil (2 t ha<sup>-1</sup>) of lime was applied as dolomite in accordance with the Fertilizer Recommendation Guide [34] and a literature review [35]. Vermicompost as organic amendments were applied at a rate of 1 g kg<sup>-1</sup> soil (2 t ha<sup>-1</sup>) and 0.5 g kg<sup>-1</sup> soil (1 t ha<sup>-1</sup>) for the first and second experiment, respectively at 10 days after germination.

**Seed sowing:** Twenty seeds of stem amaranth were broadcasted at each of the experimental pot on 25<sup>th</sup> August 2022 (For 1<sup>st</sup> pot experiment) and on 1<sup>st</sup> September 2022 (For 2<sup>nd</sup> pot experiment).

**Data collection:** Plant heights, perpendicular canopy with canopy widest width were measured three times from all plants during the growth stage. Furthermore, plant growth index was calculated by using following formula [36]:

$$\text{Plant growth index (PGI)} = \frac{\text{Plant height}}{2} + \frac{\text{Perpendicular canopy width} + \text{Canopy widest width}}{4} \quad (3)$$


At harvest, stem, leaf and root biomasses were recorded separately and they were dried in oven.

**Collection, preparation & analysis of plant samples:** While harvesting, plant samples were taken, air dried, oven dried, ground and kept in pot for analysis. The mineral nutrients in the processed plant samples under two studies were quantitatively determined in laboratory (SI 2).

**Nutrient uptake calculation:** Nutrient uptake by stem was calculated by using following formula:

$$\text{Nutrient uptake in stem (mg/kg)} = \frac{\text{Nutrient concentration in stem (\%)} \times \text{Yield (mg/pot) (dry basis)}}{100} \quad (4)$$

**Statistical analysis and graphical presentation:** Treatment effects were determined using analysis of variance (ANOVA), and treatment means were compared using the Duncan Multiple Range Test (DMRT) at a 5% level of significance in accordance with the basic principles outlined by Gomez and Gomez [37]. Obtained dataset were statistically analyzed in R console platform (38). Plant growth index (PGI) was graphically presented in line graph against different DAGs in spreadsheet.



**Fig. 1. Study sites and location of individual sediment samples (n=40 for each wetland) of the wetlands**

### 3. RESULTS

**Experiment 01:** Assessment of various wetlands (marsh) sediments in respect to their physicochemical attributes and contents of contaminants of heavy metal

**Physicochemical attributes of initial wetlands' sediments used in the study:** Physical attributes of wetlands' sediments are shown in Table 1. The overall pH value indicates that soils were very strongly to strongly acidic. The highest soil pH (5.84) and EC (0.080 ds/m) were found in wetland sediment of hilly area II while lowest pH (4.45) was found in wetland sediment of hilly area I, and the lowest EC (0.067 ds/m) was in wetland sediment of floodplain area. The overall OC was observed as low to very low (0.84-1.52 %) in studied sediments. The highest PD (2.25 g/cc) was observed in wetland sediment of floodplain area followed by wetland sediment of hilly area I.

Total N, exchangeable K, soil extractable P, S, Ca, Mg, Zn, Fe, Mn and Cu are shown in Table 2. The highest amount of total N, soil extractable S, Fe, Mn and Cu figuring 0.09%, 69.81 mg/kg, 16.15 mg/kg, 7.30 mg/kg and 1.48 mg/kg, respectively were found in the sediment of hilly area I. Exchangeable K (0.18 meq/100g soil), soil extractable P (235.15 mg/kg), Ca (2.78 meq/100g soil) and Zn (3.08 mg/kg) were observed on the highest in the wetland sediment of hilly area II. On the other hand, soil extractable Mg (1.77 meq/100g soil) was observed on higher in the sediment of floodplain area. Among the observed heavy metals, only Cd, Pb and Ni were found higher than the maximum permissible addition for all sediments (Table 3).

The highest Cd (1.14 mg/kg) was found in the wetland sediment of hilly area II followed by wetland sediment of hilly area I. The highest Pb (164.71 mg/kg) was found from wetland sediment of floodplain area followed by wetland sediment of hilly area I. Concentration of other contaminants such as Cu, Zn, and Ni were found lower than the maximum permissible addition (Table 3). But the collective risk of all heavy metals was non-dangerous for all sediments (Tables 5 & 6).

**Experiment 02:** Evaluation of sediments as media for pot culturing of a fast-growing crop stem amaranth in Eutric Gleysols and Dystic Cambisols

**Plant growth index (PGI) in Eutric Gleysols:** At 10 days after germination (DAG), the highest PGI

(9.77 cm) was observed from T<sub>4</sub> treatment receiving wetland sediment of hilly area I (Fig. 2) while the lowest one (4.25 cm) was found from T<sub>3</sub> treatment. At 20 DAG, the highest PGI (29.03 cm) was observed in T<sub>5</sub> treatment and the lowest PGI was 18.10 cm from T<sub>1</sub> (Eutric Gleysols + wetland sediment of hilly area I) treatment. At harvesting, the highest PGI (75.06 cm) was found in T<sub>5</sub> treatment (wetland sediment of hilly area II), and the lowest one (34.01 cm) was found in T<sub>1</sub> treatment.

In case of combined use of different wetlands' sediment and Eutric Gleysols as compared with T<sub>0</sub> treatment, the highest PGI (61.13 cm) was observed in T<sub>2</sub> treatment at harvest. On the other hand, different wetlands' sediment (when considered as sole utilization) compared with T<sub>0</sub> treatment, the highest PGI (75.06 cm) was observed in T<sub>5</sub> treatment at harvest.

**Plant growth index (PGI) in Dystic Cambisols:** In Dystic Cambisols, the highest PGI (13.96 cm) was observed at 10 DAG from T<sub>5</sub> treatment receiving wetland sediment of hilly area II (Fig. 3). The lowest one (6.17 cm) was found from T<sub>6</sub> treatment (wetland sediment of floodplain area). At 20 DAG, the highest PGI (34.69 cm) were observed from T<sub>5</sub> treatment receiving wetland sediment of hilly area II. The lowest PGI was 18.16 cm found from T<sub>2</sub> treatment. At harvesting, the highest PGI (65.91 cm) was observed in T<sub>5</sub> treatment and the lowest one (51.63 cm) was found in T<sub>2</sub> treatment. Almost all cases, treatment T<sub>5</sub> gave higher PGI for both studies. Although treatment T<sub>4</sub>, T<sub>5</sub> and T<sub>6</sub> were the same for both pot studies, the higher PGI was observed in 2nd pot study, might be due to temporal variation of the growth.

Combined use of sediments of Hakaluki haor and Dystic Cambisols exhibited highest PGI (58.57 cm) at the time of harvest. Conversely, as compared to the T<sub>0</sub> treatment, the highest PGI (65.91 cm) was observed in the T<sub>5</sub> treatment at harvest.

**Fresh biomass production:** Fresh biomass production in different plant parts of stem amaranth grown in both pot experiments varied significantly due to the application of different treatments (Table 5). In the 1<sup>st</sup> pot experiment, the highest fresh weights of stem, root and leaf were recorded as 148.52, 25.27 and 72.71 g pot<sup>-1</sup>, respectively in the T<sub>5</sub> treated pot. And, the minimum weights of stem, root and leaf were 29.26, 5.32 and 14.63 g pot<sup>-1</sup>, respectively in the pot with the treatment T<sub>1</sub>. Consequently, the T<sub>5</sub>

**Table 1. Physical attributes of three wetlands' sediments and farmer's field soils**

Sediment / Soil	pH	OC (%)	EC (ds/m)	PD (g/cc)	Texture	Consistency
Wetland sediment of hilly area I	4.45	1.52	0.067	2.16	Sandy clay loam	Firm
Wetland sediment of hilly area II	5.84	0.84	0.080	2.23	Sandy clay loam	Very firm
Wetland sediment of floodplain area	5.34	1.04	0.063	2.25	Sandy loam	Firm
Eutric Gleysols	6	1.48	0.082	2.39	Clay loam	Very firm
Dystic Cambisols	6.1	1.00	0.084	2.23	Sandy loam	Firm

OC = Organic carbon, EC = Electrical conductivity, PD = Particle density

**Table 2. Chemical attributes of initial sediments and farmer's field soils**

Elements	Wetlands' Sediments			Farmers' Field Soils	
	Hilly area I	Hilly area II	Floodplain area	Eutric Gleysols	Dystic Cambisols
N <sup>T</sup> (%)	0.09 (Very low)	0.07 (Very low)	0.06 (Very low)	0.095 (Low)	0.07 (Very low)
K <sup>Exc</sup> (meq/100g soil)	0.13 (Low)	0.18 (Medium)	0.05 (Very low)	0.08 (Very low)	0.09 (Low)
P <sup>E</sup> (mg/kg)	40.62 (Very high)	235.15 (Very high)	64.13 (Very high)	11.1 (Medium)	5.00 (Medium)
S <sup>E</sup> (mg/kg)	69.81 (Very high)	57.74 (Very high)	29.17 (Very high)	10.5 (Low)	9.80 (Low)
Ca <sup>E</sup> (meq/100g soil)	1.59 (Low)	2.78 (Low)	2.15 (Low)	-	-
Mg <sup>E</sup> (meq/100g soil)	1.50 (Optimum)	1.52 (Optimum)	1.77 (High)	-	-
Zn <sup>E</sup> (mg/kg)	1.85 (High)	3.08 (Very high)	2.63 (Very high)	-	-
Fe <sup>E</sup> (mg/kg)	16.15 (Very high)	11.50 (High)	9.83 (High)	-	-
Mn <sup>E</sup> (mg/kg)	7.30 (Very high)	2.67 (Optimum)	4.44 (Very high)	-	-
Cu <sup>E</sup> (mg/kg)	1.48 (Very high)	0.85 (Very high)	1.25 (Very high)	-	-

T= Total, Exc= Exchangeable, E= Extractable; Interpretation (within the parenthesis) for the soil test values was done based on FRG (2018)

**Table 3. Nonspecific biomass targeted soil (sediment) pollution risk by heavy metals**

Elements	Observed analytical value (mg/kg)			MPA as Reference value (RV) (mg/kg)	Ratio (SH-I /RV)	Ratio (SH-II/RV)	Ratio (SF/RV)	Collective Risk of All heavy metals and comments (As adapted from Saet et al. [33] (Zc)		
	Sediment of Hilly area I (SH-I)	Sediment of Hilly area II (SH-II)	Sediment of Floodplain area (SF)					SH-I	SH-II	SF
<b>Cu (mg/kg)</b>	1.48	0.85	1.25	3.5	0.42	0.24	0.36		15.59: No	13.71: No
<b>Zn (mg/kg)</b>	1.85	3.08	2.63	16	0.12	0.19	0.16	16.93: No	Dangerous	Dangerous
<b>Cd (mg/kg)</b>	1.06	1.14	1.05	0.76	1.39	1.5	1.38	Dangerous		
<b>Pd (mg/kg)</b>	102.88	64.71	164.71	55	1.87	1.18	2.99			
<b>Ni (mg/kg)</b>	44.54	42.86	33.33	2.6	17.13	16.48	12.82			

MPA= Maximal permissible addition

**Table 4. Plant specific (stem amaranth) targeted soil (sediment) pollution risk by heavy metals used as plant nutrients**

Elements	Current Status			Risk of Heavy Metals as Nutrient Element (Zc <sub>n</sub> )			Zc <sub>n</sub> values with annotation			
	SH-I	SH-II	SF	SH-I	SH-II	SF				
<b>Cu (mg/kg)</b>	1.48	0.85	1.25	0.88	0.25	0.65	Less than zero (<0): Lower than plant requirement	Equal or near to zero (0): Optimum	Greater than zero (>0): High to very high	Great greater than zero (>>0): Excess to toxic
<b>Zn (mg/kg)</b>	1.85	3.08	2.63	0.05	1.58	1.13				

SH-I= Sediment of Hilly area I, SH-II= Sediment of Hilly area II, SF= Sediment of Floodplain area

**Table 5. Fresh biomass of stem amaranth produced in first and second pot experiments under different treatments at harvest**

Treatments	Fresh biomass (g pot <sup>-1</sup> )							
	1st Pot Experiment (Eutric Gleysols)				2nd Pot Experiment (Dystic Cambisols)			
	Stem weight	Root weight	Leaf weight	Total Fresh Weight	Stem weight	Root weight	Leaf weight	Total Fresh Weight
T <sub>0</sub>	34.58 ± 1.15 ef	6.65 ± 0.58 ef	19.06 ± 1.20 e	60.29 ± 0.66 e	67.39 ± 1.66 e	9.31 ± 0.58 e	31.92 ± 1.73 f	108.62 ± 3.71 e
T <sub>1</sub>	29.26 ± 2.51 f	5.32 ± 0.58 f	14.63 ± 1.15 e	49.21 ± 3.60 e	85.56 ± 2.60 d	11.97 ± 0.58 de	23.50 ± 1.45 g	121.03 ± 3.51 e
T <sub>2</sub>	106.4 ± 4.16 b	17.73 ± 0.88 b	55.42 ± 0.88 b	179.55 ± 3.79 b	59.85 ± 1.53 e	5.32 ± 0.58 f	41.67 ± 1.45 e	106.84 ± 2.73 e
T <sub>3</sub>	70.05 ± 2.02 c	13.74 ± 0.88 c	42.65 ± 2.08 c	126.35 ± 3.79 c	105.51 ± 2.96 c	16.85 ± 0.66 c	61.18 ± 3.21 c	183.54 ± 6.08 c
T <sub>4</sub>	53.64 ± 2.33 d	11.08 ± 0.33 cd	32.36 ± 1.45 d	97.09 ± 2.88 d	91.77 ± 2.30 cd	15.07 ± 0.88 cd	50.54 ± 1.15 d	157.38 ± 2.33 d
T <sub>5</sub>	148.52 ± 4.33 a	25.27 ± 1.53 a	72.71 ± 2.19 a	246.49 ± 7.68 a	203.93 ± 7.51 a	22.61 ± 1.00 a	100.19 ± 2.30 a	330.73 ± 9.26 a
T <sub>6</sub>	43.00 ± 2.02 de	9.75 ± 0.33 de	29.70 ± 1.20 d	82.46 ± 3.21 d	131.23 ± 4.63 b	26.6 ± 1.53 b	70.49 ± 2.19 b	224.33 ± 4.06 b
CV (%)	9.51	14.87	9.21	7.93	8.33	13.33	8.64	6.60
LS	***	***	***	***	***	***	***	***

Values are mean ± SEM (Standard Errors of Mean), Means followed by different letter in a column are significantly different at 0.1% level by LSD, LS = Level of significance, CV = Co-efficient of variation, T<sub>0</sub>: Farmer's field soil (FFS) (Eutric Glaysols for 1st pot experiment and Dystic Cambisols for 2nd pot experiment); T<sub>1</sub>: FFS + Wetland sediment of hilly area I (1:1); T<sub>2</sub>: FFS + Wetland sediment of hilly area II (1:1); T<sub>3</sub>: FFS + Wetland sediment of floodplain area (1:1); T<sub>4</sub>: Wetland sediment of hilly area I; T<sub>5</sub>: Wetland sediment of hilly area II; T<sub>6</sub>: Wetland sediment of floodplain area.

**Table 6. Dry biomass of stem amaranth produced in first and second pot experiments under different treatments**

Treatments	Dry biomass (g pot <sup>-1</sup> )							
	1st Pot Experiment (Eutric Gleysols)				2nd Pot Experiment (Dystic Cambisols)			
	Stem weight	Root weight	Leaf weight	Total Dry Weight	Stem weight	Root weight	Leaf weight	Total Dry Weight
T <sub>0</sub>	3.43 ± 0.08 d	0.83 ± 0.01 ef	3.47 ± 0.38 d	7.74 ± 0.44 f	4.03 ± 0.11 f	0.79 ± 0.03 e	4.02 ± 0.08 c	8.80 ± 0.06 f
T <sub>1</sub>	1.80 ± 0.14 e	0.60 ± 0.02 f	4.02 ± 0.19 cd	6.42 ± 0.04 f	5.09 ± 0.10 e	1.19 ± 0.04 d	3.83 ± 0.16 c	10.11 ± 0.26 e
T <sub>2</sub>	8.41 ± 0.17 b	2.09 ± 0.02 b	5.45 ± 0.16 ab	15.96 ± 0.36 b	2.72 ± 0.05 g	0.60 ± 0.03 e	3.87 ± 0.09 c	7.18 ± 0.16 g
T <sub>3</sub>	7.55 ± 0.28 b	1.37 ± 0.06 c	5.12 ± 0.22 abc	14.05 ± 0.26 c	7.29 ± 0.28 c	1.92 ± 0.10 c	5.26 ± 0.26 b	14.67 ± 0.37 c
T <sub>4</sub>	6.38 ± 0.15 c	1.14 ± 0.02 cd	4.73 ± 0.52 bc	12.24 ± 0.39 d	6.50 ± 0.06 d	1.37 ± 0.03 d	5.19 ± 0.29 b	13.07 ± 0.35 d
T <sub>5</sub>	9.89 ± 0.19 a	3.08 ± 0.15 a	6.06 ± 0.03 a	19.03 ± 0.06 a	10.17 ± 0.19 a	3.46 ± 0.07 a	6.58 ± 0.12 a	20.22 ± 0.11 a
T <sub>6</sub>	3.93 ± 0.36 d	0.93 ± 0.01 de	4.25 ± 0.12 cd	9.11 ± 0.48 e	8.74 ± 0.05 b	2.48 ± 0.06 b	5.95 ± 0.14 ab	17.18 ± 0.27 b
CV (%)	8.41	10.20	13.51	6.38	5.24	8.49	8.50	4.45
LS	***	***	**	***	***	***	***	***

Values are mean ± SEM (Standard Errors of Mean), Means followed by different letter in a column are significantly different at 0.1% level by LSD, LS = Level of significance, CV = Co-efficient of variation, T<sub>0</sub>: Farmer's field soil (FFS) (Eutric Glaysols for 1st pot experiment and Dystic Cambisols 2nd pot experiment); T<sub>1</sub>: FFS + Wetland sediment of hilly area I (1:1); T<sub>2</sub>: FFS + Wetland sediment of hilly area II (1:1); T<sub>3</sub>: FFS + Wetland sediment of floodplain area (1:1); T<sub>4</sub>: Wetland sediment of hilly area I; T<sub>5</sub>: Wetland sediment of hilly area II; T<sub>6</sub>: Wetland sediment of floodplain area



treatment yielded the highest fresh weight (total) of 246.49 g pot<sup>-1</sup> and the pot with T<sub>1</sub> treatment gave the lowest fresh weight (total) of 49.21 g. Treatment T<sub>5</sub> was followed by treatment T<sub>2</sub> which was followed by treatments T<sub>3</sub> in yielding total fresh biomass. In the case of utilizing a combination of sediment from various wetlands' and Eutric Gleysols compared to the T<sub>0</sub> treatment, the highest total fresh biomass (179.55 g pot<sup>-1</sup>) was recorded in T<sub>2</sub> treatment. On the other hand, as compared to the T<sub>0</sub> treatment, the highest total fresh biomass (246.49 g pot<sup>-1</sup>) was observed in the T<sub>5</sub> treatment.

Similarly, in the 2<sup>nd</sup> pot experiment, the highest fresh weight of stem, root and leaf was recorded as 203.93, 22.61 and 100.19 g pot<sup>-1</sup>, respectively in the T<sub>5</sub> treated pots. And, the lowest weights of stem and root were 59.85 and 5.32 g pot<sup>-1</sup>, respectively in the pot with the treatment T<sub>2</sub> while the lowest weight of leaf was 23.50 g pot<sup>-1</sup> in the pot with the treatment T<sub>1</sub>. Eventually, the T<sub>5</sub> treatment gave total fresh weight (highest) of 330.73 g and the T<sub>2</sub> treatment yielded the total fresh weight (lowest) of 106.84 g. Treatment T<sub>5</sub> was followed by treatment T<sub>6</sub> which was followed by treatments T<sub>3</sub> in yielding total fresh biomass. In case of integrated use of field soil and sediment, T<sub>3</sub> showed the highest total fresh biomass (183.54 g pot<sup>-1</sup>). But, the highest total fresh biomass (330.73 g pot<sup>-1</sup>) was observed in the T<sub>5</sub> treatment (Tanguar haor) among all sole use of sediments.

**Dry biomass production:** Dry biomass for different parts of stem amaranth varied significantly by the application of different treatments in both pot experiments as shown in Table 6. In case of 1<sup>st</sup> pot experiment (in Eutric Gleysols), the highest dry stem, root and leaf weight was found as 9.89, 3.08 and 6.06 g pot<sup>-1</sup>, respectively in the pot receiving T<sub>5</sub> treatment. The lowest stem and root weights were recorded as 1.80 and 0.60 g pot<sup>-1</sup>, respectively in the T<sub>1</sub> treated pot, while the lowest leaf weight was 3.47 g pot<sup>-1</sup> found from T<sub>0</sub> treatment. As a consequence, the T<sub>5</sub> treated pot gave the highest total dry weight of 19.03 g pot<sup>-1</sup> while the T<sub>1</sub> pot gave the lowest total dry weight of 6.42 g pot<sup>-1</sup>.

In case of using a combination of sediments from various wetlands and Eutric Gleysols, the highest total dry biomass (15.96 g pot<sup>-1</sup>) was recorded in the T<sub>2</sub> treatment. On the contrary, when we used sediment (sole) from different wetlands the highest total dry biomass (19.03 g pot<sup>-1</sup>) was observed in the T<sub>5</sub> treatment.

In the 2<sup>nd</sup> pot experiment (in Dystic Cambisols), the highest dry stem, root and leaf weights were found as 10.17, 3.46 and 6.58 g pot<sup>-1</sup>, respectively in the pot receiving T<sub>5</sub> treatment. The lowest stem and root weights were recorded as 2.72 and 0.60 g pot<sup>-1</sup>, respectively in the T<sub>2</sub> treated pot, while the lowest leaf weight was 3.83 g pot<sup>-1</sup> found from T<sub>1</sub> treatment. As a consequence, the T<sub>5</sub> treated pot gave the highest total dry weight of 20.22 g pot<sup>-1</sup> while T<sub>2</sub> treated pot gave the lowest total dry weight of 7.18 g pot<sup>-1</sup>. In case of using a combination of sediment from various wetlands and Dystic Cambisols, the highest total dry biomass (14.67 g pot<sup>-1</sup>) was recorded in the T<sub>2</sub> treatment. On the contrary, considering sole use of sediments from different wetlands, the highest total dry biomass (20.22 g pot<sup>-1</sup>) was observed in the T<sub>5</sub> treatment.

**Nutrients uptake in stem:** In case of first pot experiment, significant differences were observed for nutrient uptake in stem due to the different treatments which are shown in Fig. 4. Use of wetland sediment of hilly area II as a growing media (T<sub>5</sub>) resulted the highest P, K, S, Ca, Mg, B, Zn, Fe and Mn uptakes figuring 2399.98, 11915.40, 3087.86, 5733.04, 1073.38, 114.50, 130.75, 801.96 and 156.63 mg kg<sup>-1</sup> respectively. But highest Cu (9.89 mg kg<sup>-1</sup>) uptake was recorded from combined use of Eutric Gleysols and wetland sediment of hilly area II (T<sub>2</sub>).

When considering the mixed treatment in comparison to the control (T<sub>0</sub>), the uptake of all mineral nutrients was found to be highest in the T<sub>2</sub> treatment compared to the other treatments. The lowest uptake of P, K, S, Ca, Mg, B, Zn, Fe, Mn and Cu were in pots with combined use of Eutric Gleysols and wetland sediment of hilly area I (T<sub>1</sub>).

In case of second pot experiment, significant differences were observed for nutrient uptake in stem due to the different treatments which are shown in Fig. 4. The highest uptake of P, K, Ca and Mn figuring 3955.86, 12035.90, 3341.68 and 165.46 mg kg<sup>-1</sup> respectively, were observed in pots with the use of wetland sediment of hilly area II as a growing media (T<sub>5</sub>). But the highest S, Mg, Zn, Fe and B uptakes figuring 2519.89, 1043.86, 81.71, 1753.15 and 368.36 mg kg<sup>-1</sup> respectively, were recorded in wetland sediment of floodplain area (T<sub>6</sub>).

In case of integrated use of sediments and fields soils, the uptake of all mineral nutrients was

found to be highest in the T<sub>3</sub> treatment compared to the other treatments. The lowest uptakes of P, S, Ca, Mg, B, Zn and Mn in pots with the combined use of Dyctic Cambisols and wetland sediment of hilly area II (T<sub>1</sub>). On the other hand, the lowest uptakes of K, Fe and Cu were in pots containing Dyctic Cambisols (T<sub>0</sub>).

#### 4. DISCUSSION

**Physical attributes of different wetland sediments:** The pH value of the studied sediments was observed as strongly to very strongly acidic might be due to the presence of acidic ions. Due to residence in submerged condition, it was supposed that sediments in wetland were in neutral or near neutral in reaction in *in-situ* condition. Similarly, soil OC was observed low to very low. In actual submerged condition the OC were expected to be higher. But, after soil sample collection, these soils got oxidized during drying of wet

sediments and eventually OM content lowered down.

**Soil (sediment) pollution risk by heavy metals:** The heavy metal concentrations of the different sediments are shown in Table 3. For nonspecific biomass, overall collective risk of all heavy metals of the studied heavy metals in the study area could be regarded as not dangerous as per the literature (Z<sub>c</sub>) by Saet et al. [33]. It may be due to the concentrations of heavy metals in sediments were lower than the maximal permissible addition (MPA), (MPA as reported by Crommentuijn et al. [39]. On the contrary, in case of plant specific (stem amaranth) targeted soil (sediment) pollution risk by heavy metals used as plant nutrients (Cu and Zn) were observed on very high to excess in the studied area as per our proposed pollution risk index (Z<sub>c<sub>n</sub></sub>) (Table 4). Baran et al. [40] also confirmed that the addition of sediment in a dose of 50% to a sandy soil increased the contents of Zn and Cu, which supports our finding.




Fig. 2. Plant growth index (PGI) in Eutric Gleysols at different days after germination (DAG)




Fig. 3. Plant growth index (PGI) in Dyctic Cambisols at different days after germination (DAG)




Fig. 4. Nutrients uptake by stem amaranth

**Growth, yield and nutrient uptake of stem amaranth:** The application of different treatments considerably changed the growth and yield parameter of stem amaranth. For both net house studies, in all cases, the T<sub>5</sub> treatment containing wetland sediment of hilly area II performed better than the remaining treatments in contributing increment of growth and yield parameters. This result might be due to the plant's ability to take up significantly greater amounts of P from the sediment than from sole soil, as wetland sediment of hilly area II contained more easily soluble P than sole farmers' soil (Table 2). The statement of Baran et al. [40] supports this finding, who states that different doses of sediments to sandy soil significantly increased the content of available P. In addition to P, the majority of easily soluble macro- and -micronutrients were considerably more abundant in the sediment than the others, which enhanced suitable conditions (fertility) for plant growth. These nutrients are likely originated from upper agricultural soils, which, through erosion, and deposition reached the bottom of the lake sediment [41,42,43]. Canet et al. [21] also reported an increase in lettuce yield due to the nutrient contents of the sediments and the possible improvement of the cation exchange capacity. Our finding is supported by recent studies reporting that the sediment mixture had comparable growth performance for holm oak seedlings [44], ornamental Red Robin photinia [45], lettuce [21], and strawberry [19] compared to others. Compared to the 1st study, 2nd study showed the higher growth and yield in almost all cases. This was due to, as per our observation, the fact that seedlings under 1st pot experiment faced heat stress. And, 2nd pot experiment escaped the period for late sowing.

The uptake of nutrients such as P, K, S, Ca, Mg, B, Zn, Fe and Mn (for 1st pot experiment) and P, K, S and Ca (for 2nd pot experiment) in stem were found to be higher in the pot with treatment T<sub>5</sub> using wetland sediment of hilly area II. This could be due to the higher initial nutrients' concentration in this sediment. Our findings partially align with Islam et al. [46] who observed that pond sediments revealed a higher quantity of nutritional contents in Indian spinach.

#### 4. CONCLUSIONS

The collective risk of all heavy metals was non-dangerous for all studied sediments collected from wetland sediment of hilly area I, wetland sediment of hilly area II, and wetland sediment of

a floodplain area. Wetland sediment of hilly area II as a vegetable crop growing media gave better growth and yield among the studied sediments. So, the use of wetland sediment as a growing medium can be beneficial to produce amaranth as vegetable especially for flash flood prone wetland areas. Further field trial is needed to extrapolate the findings in a larger scale.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### FUNDING

The Ministry of Science and Technology (MoST) of the Government of the People's Republic of Bangladesh for granting funds to conduct the research.

#### DATA AVAILABILITY

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

#### ACKNOWLEDGEMENTS

We would like to acknowledge the Ministry of Science and Technology (MoST) of the Government of the People's Republic of Bangladesh for granting funds to conduct the research.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Wikipedia: About 'wetlands'. Retrieved April 09, 2024 from Wikipedia: <https://en.wikipedia.org/wiki/Wetland>.
2. <https://www.epa.gov/wetlands/what-wetland>.
3. Renella G. Recycling and reuse of sediments in agriculture: Where is the problem? Sustainability. 2021;13(4):1648.
4. De Araújo JC, Güntner A, Bronstert A. Loss of reservoir volume by sediment deposition and its impact on water

- availability in semiarid Brazil. *Hydrological Sciences Journal*. 2006;51(1):157-170.
5. Owens PN, Batalla RJ, Collins AJ, Gomez B, Hicks DM, Horowitz AJ, Trustrum NA. Fine-grained sediment in river systems: Environmental significance and management issues. *River Research and Applications*. 2005;21(7):693-717.
  6. Yozzo DJ, Wilber P, Will RJ. Beneficial use of dredged material for habitat creation, enhancement, and restoration in New York–New Jersey Harbor. *Journal of Environmental Management*. 2004;73(1):39-52.
  7. Kondolf GM, Gao Y, Annandale GW, Morris GL, Jiang E, Zhang J, Yang CT. Sustainable sediment management in reservoirs and regulated rivers: Experiences from five continents. *Earth's Future*. 2014;2(5):256-280.
  8. Wójcikowska-Kapusta A, Smal H, Ligęza S. Contents of selected macronutrients in bottom sediments of two water reservoirs and assessment of their suitability for natural use. *Journal of Water and Land Development*. 2018;38:147-153.
  9. Mymrin V, Stella JC, Scremim CB, Pan RC, Sanches FG, Alekseev K, Fortini OM. Utilization of sediments dredged from marine ports as a principal component of composite material. *Journal of Cleaner Production*. 2017;142:4041-4049.
  10. SedNet. Contaminated sediments in European river basins. *European Sediment Research Network*; 2014.
  11. Szara-Bąk M, Baran A, Klimkowicz-Pawlas A. Recycling of bottom sediment to agriculture: Effects on plant growth and soil properties. *Journal of Soils and Sediments*. 2023;23(1):539-551.
  12. Baran A, Jasiewicz C, Tarnawski M. Effect of bottom sediment supplement to light soil on the content and uptake of macro elements by maize. *Ecological Chemistry and Engineering. A*. 2012;19(8):863-871.
  13. Jasiewicz C, Madeyski M, Tarnawski M, Baran A. Effect of bottom sediment supplement to soil on yield and chemical composition of maize. *Ecological Chemistry and Engineering. A*. 2011;18(11):1505-1514.
  14. Kiani M, Raave H, Simojoki A, Tammeorg O, Tammeorg P. Recycling lake sediment to agriculture: Effects on plant growth, nutrient availability, and leaching. *Science of the Total Environment*. 2021;753:141984.
  15. Wiśniowska-Kielian B, Niemiec M. Effect of bottom sediment addition to the substratum on the quality of produced maize (*Zea Mays L.*) biomass. *Chemia i Inżynieria Ekologiczna*. 2007a;14(5-6):581-589.
  16. Wiśniowska-Kielian B, Niemiec M. Effect of increasing share of bottom sediment in the substratum on cadmium and lead uptake by the plants. *Chemia i Inżynieria Ekologiczna*. 2007b;14(5-6):591-599.
  17. Tarnawski M, Baran A, Koniarz T. The effect of bottom sediment supplement on changes of soil properties and on the chemical composition of plants. *Geology, Geophysics and Environment*. 2015;41(3):285-285.
  18. Macía P, Fernández-Costas C, Rodríguez E, Sieiro P, Pazos M, Sanromán MA. Technosols as a novel valorization strategy for an ecological management of dredged marine sediments. *Ecological Engineering*. 2014;67:182-189.
  19. Tozzi F, Del Bubba M, Petrucci WA, Pecchioli S, Macci C, García FH, Giordani E. Use of a remediated dredged marine sediment as a substrate for food crop cultivation: Sediment characterization and assessment of fruit safety and quality using strawberry (*Fragaria x ananassa* Duch.) as model species of contamination transfer. *Chemosphere*. 2020;238:124651.
  20. Wobus A, Bleul C, Maassen S, Scheerer C, Schuppler M, Jacobs E, Röske I. Microbial diversity and functional characterization of sediments from reservoirs of different trophic state. *FEMS Microbiology Ecology*. 2003;46(3):331-347.
  21. Canet R, Chaves C, Pomares F, Albiach R. Agricultural use of sediments from the Albufera Lake (Eastern Spain). *Agriculture, Ecosystems and Environment*. 2003;95(1):29-36.
  22. Kazberuk W, Szulc W, Rutkowska B. Use bottom sediment to agriculture—??Effect on plant and heavy metal content in soil. *Agronomy*. 2021;11(6):1077.
  23. Braga BB, Junior FN, Barbosa RM, Brito POB, Martins K, Medeiros PHA, Gondim FA. Biomass production and antioxidative enzyme activities of sunflower plants growing in substrates containing sediment from a tropical reservoir. *Journal of Agricultural Science*. 2017;9(5):1-12.
  24. Baran A, Tarnawski M, Koniarz T, Jasiewicz C. Agricultural use of sediments from Narożniki reservoir—yield and

- concentration of macronutrients and trace elements in the plant. *Infrastrukturai Ekologia Terenów Wiejskich*; 2016.
25. Capra GF, Grilli E, Macci C, Vacca S, Masciandaro G, Ceccanti B, Buondonno A. Lake-dredged material (LDM) in pedotechnique for the restoration of Mediterranean soils affected by erosion/entisolization processes. *Journal of Soils and Sediments*. 2015;15:32-46.
  26. Nygård A, Purhonen M. Hollolan Työtjärven pohjasedimentin hyötykäyttö; 2019.
  27. Sapota G, Dembska G, Bogdaniuk M, Holm G. Environmental policy and legislation on dredged material in the Baltic Sea region—???analysis. In 2012 IEEE/OES Baltic International Symposium (BALTIC). IEEE. 2012;1-15.
  28. Finnish Water Directive; 2011. Available:<https://www.finlex.fi/fi/laki/smur/2011/20110587>
  29. Ferrans L, Jani Y, Gao L, Hogland W. Characterization of dredged sediments: A first guide to define potentially valuable compounds—the case of Malmfjärden Bay, Sweden. *Advances in Geosciences*. 2019;49:137-147.
  30. Couvidat J, Chatain V, Bouzahzah H, Benzaazoua M. Characterization of how contaminants arise in a dredged marine sediment and analysis of the effect of natural weathering. *Science of the Total Environment*. 2018;624:323-332.
  31. Talukder N, Murshed S (2021) [www.banglapidia.org](http://www.banglapidia.org).
  32. Alam S, Hossain MS (2021) [www.banglapidia.org](http://www.banglapidia.org).
  33. Saet YE, Revich BA, Yanin EP. *Geochemistry of the environment*. Nedra, Moscow. 1990;82.
  34. FRG. Fertilizer Recommendation Guide, Bangladesh Agriculture Research Council (BARC). Farmgate Dhaka. 2018;1215: 274.
  35. Nazrul MI, Shaheb MR. Performance of French bean (*Phaseolus vulgaris* L.) genotypes in Sylhet region of Bangladesh. *Bangladesh Agronomy Journal*. 2016; 19(1):37-44.
  36. Liu R, Gu M, Huang L, Yu F, Jung SK, Choi HS. Effect of pine wood biochar mixed with two types of compost on growth of bell pepper (*Capsicum annuum* L.). *Horticulture, Environment, and Biotechnology*. 2019;60:313-319.
  37. Gomez KA, Gomez AA. *Statistical procedures for agricultural research*. John Wiley and Sons; 1984.
  38. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria; 2021.
  39. Crommentuijn T, Polder MD. Maximum permissible concentrations and negligible concentrations for metals, taking background concentrations into account; 1997.
  40. Baran A, Tarnawski M, Urbaniak M. An assessment of bottom sediment as a source of plant nutrients and an agent for improving soil properties. *Environmental Engineering and Management Journal*. (Some of the journal names are in italics. Journal names should either be written in italics or normally). 2019;18(8):1647-1656
  41. Kusic I, Basic F, Nestroy O, Mesic M, Butorac A. Chemical properties of eroded soil material. *Journal of Agronomy and Crop Science*. 2002;188(5):323-334.
  42. Fonseca RM, Barriga FJ, Conceição PI. Clay minerals in sediments of Portuguese reservoirs and their significance as weathering products from over-eroded soils: A comparative study of the Maranhão, Monte Novo and Divor Reservoirs (South Portugal). *International Journal of Earth Sciences*. 2010;99:1899-1916.
  43. De Vincenzo A, Covelli C, Molino AJ, Pannone M, Ciccaglione M, Molino B. Long-term management policies of reservoirs: Possible re-use of dredged sediments for coastal nourishment. *Water*. 2018;11(1):15.
  44. Ugolini F, Mariotti B, Maltoni A, Tani A, Salbitano F, Izquierdo CG, Tognetti R. A tree from waste: Decontaminated dredged sediments for growing forest tree seedlings. *Journal of Environmental Management*. 2018;211:269-277.
  45. Mattei P, Pastorelli R, Rami G, Mocali S, Giagnoni L, Gonnelli C, Renella G. Evaluation of dredged sediment co-composted with green waste as plant growing media assessed by ecotoxicological tests, plant growth and microbial community structure. *Journal of Hazardous Materials*. 2017;333:144-153.
  46. Islam MA, Jahan N, Quadir QF, Haque SM. Assessment of fish pond sediments for growth, yield and nutritional quality of Indian spinach (*Basella alba* L.). *Archives*

- of Agriculture and Environmental Science. 2020;5(1):33-39.
47. Jackson ML. Soil chemical analysis. New Delhi: Prentice-Hall of India Private Limited; 1973.
  48. Bouyoucos GJ. Hydrometer method improved for making particle size analyses of soils 1. Agronomy Journal. 1962;54(5): 464-465.
  49. Food and Agriculture Organization of the United Nations (FAO), (2015). World Fertilizer Trends and Outlook. 2018;1–66.
  50. Nelson DW, Sommers LE. Total carbon, organic carbon, and organic matter. Methods of Soil Analysis: Part 3 Chemical Methods. 1996;5:961-1010.
  51. Bremner JM, Mulvaney CS. Total Nitrogen. In: Page AL, Miller RH and Keeney DR (editors), Methods of Soil Analysis, Part 2, Chemical and Microbiological Properties, American Society of Agronomy and Soil Science Society of America Inc. Madison Wisconsin USA 595-624; 1982.
  52. Bray RH, Kurtz LT. Determination of total, organic, and available forms of phosphorus in soils. Soil Science. 1945;59(1):39-46.
  53. Fox RL, Olson RA, Rhoades HF. Evaluating the sulfur status of soils by plant and soil tests. Soil Science Society of America Journal. 1964;28(2):243-246.
  54. Lindsay WL, Norvell W. Development of a DTPA soil test for zinc, Iron, manganese, and copper. Soil Science Society of America Journal. 1978;42(3):421-428.
  55. Yoshida S, Forno AD, Cock JA, Gomez KA. Physiological studies of rice. 2nd Edition. International Rice Research Institute, Manila, Philippines; 1976.
  56. Chapman HD, Pratt PF. Methods of analysis for soils, plants and waters. Division of Agricultural Sciences. University of California, Berkeley. 1961; 169-176.

### SI 1 Methods used for the analysis of soil (sediment) physicochemical parameters

Parameters	Analytical methods
Soil pH	The pH of the soil was measured using a glass-electrode pH meter while maintaining a 1:2.5 soil-water ratio [47].
Soil texture (%)	Particle analysis was done by hydrometer method [48].
Soil consistency	Moist soil consistency test was done by field test of moist soil [49].
Soil color	Soil color test was done by using Munsell color chart.
Particle density	Particle density was determined by volumetric flask method as described by Nelson and Sommers [50].
Electrical conductivity (dS/m)	Electrical conductivity (Soil water ratio 1:5) was determined by Metrohm EC Meter.
Organic carbon (%)	Determined by wet oxidation method [50].
Total nitrogen (%)	Determined by Kjeldahl method [51].
Extractable phosphorus (mg/kg)	Extracted from the soil by shaking with ammonium fluoride extracting solution following Bray and Kurtz method [52].
Exchangeable potassium (meq/100 g soil)	Extracted with 1.0 N NH <sub>4</sub> OAc (pH 7) and determined by flame photometry [47].
Extractable sulphur (mg/kg)	Extracted the soil samples with CaCl <sub>2</sub> solution (0.15 %) and determined by turbidimetric method [53].
Extractable Zn, Fe, Mn, Cu, Cd, Ni, Cr (mg/kg)	Determined by DTPA extraction method [54].
Extractable Ca (meq/100 g soil)	Extracted with 1.0 N NH <sub>4</sub> OAc (pH 7) and determined by flame photometry [47].

### SI 2 The analytical methods followed to determine the parameters of plant samples.

Elements	Analytical methods
Phosphorus	Samples were digestion with nitric acid and determined by Spectrophotometry [55].
Potassium	Samples were digestion with nitric acid and determined by flame photometry [55].
Sulphur	Samples were digestion with nitric and perchloric acid. Determined by precipitation as barium sulphate [56].
Calcium, Magnesium, Zinc, Iron, Manganese, Copper and Boron	Samples were digestion with nitric acid and determined by atomic absorption spectrophotometry [55].

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/120669>