



## **A Review on *In-vitro* Haploid Production in Orchids**

**Teena Jaswal<sup>a</sup> and Saranjeet Kaur<sup>a\*</sup>**

<sup>a</sup> Department of Biosciences, University Institute of Biotechnology, Chandigarh University, District-Mohali, Punjab, India.

### **Authors' contributions**

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

### **Article Information**

DOI: 10.9734/JPRI/2021/v33i57B34023

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/77563>

**Review Article**

**Received 06 October 2021**  
**Accepted 12 December 2021**  
**Published 14 December 2021**

## **ABSTRACT**

The aim of writing this paper is to review production of haploids in orchids *in vitro*. Haploids possess half number of chromosomes and do not undergo fertilization. *In vitro* conditions provide necessary nutrients and conditions that are required for growth of haploid plantlets. The natural breeding cycle of orchids is very slow as well as unpredictable. To reduce this time, the technique of *in vitro* haploid production is used. *In vitro* conditions can decrease or shorten the time required for juvenile period in Orchids. Haploid plants are produced from *in vitro* haploid culture. This technique is useful to produce homozygous pure lines and to increase the yield of a plant.

**Keywords:** *In vitro*; haploid production; orchidaceae; orchid seeds; angiosperms; monocots.

## **1. INTRODUCTION**

Orchids are the most favourite as well as the most expensive cut-flowers. They are commercially very valuable. Due to the high market value of Orchids, there would be pressure on the cultivators to produce a large number of Orchids. The only hurdle in this process is the natural breeding program of Orchids. Family Orchidaceae contains very beautiful and vibrant

flowers. Orchids are found almost all over the world, except in the freezing polar regions and extremely hot deserts. Orchids are mostly found in tropical forests [1]. The flowers of the Orchidaceae family are of different shapes and radiant colours [2]. Orchidaceae is a very interesting family because of the colours of its flowers and the exceptional structures of the flowers [1]. These flowers are commercially very useful. Orchids flowers have high artistic value.

\*Corresponding author: E-mail: [sksekhkone7632@cumail.in](mailto:sksekhkone7632@cumail.in);

Orchids are endowed with economic worth at par with other ornamentals [3].

Orchidaceae contains roughly 35,000 species, in 1,000 genera and 1,00,000 hybrids [4]. Orchids are around 8% to 10% of all flowering plants [4]. The annual orchid flowers production in India is approximately 1000 tonnes making a small share of 0.01% in international market [2,5]. Orchids are the most expensive cut- flowers. Besides their ornamental value, some genera also provide food products and medicines [4]. The commercial value for orchid flowers is very high, due to which they are in high demand all over the world. The commercial demand for orchid plants is increasing day by day; so, the improvements in orchid cultivation and their conservation can effectively develop the economy of our country.

## **2. *In-vitro* ANTHOR, MICROSPOROVARY AND OVULE CULTURE**

Growing orchids naturally is a time-consuming and slow process [6]. Orchids have a long vegetative phase before they flower [7]. The breeding cycle of orchidaceae is also very slow and unpredictable [8]. To reduce or minimize this time period, the technique of *in vitro* haploid production is used. Haploids are useful in the shortening of the breeding cycle in orchids [9]. This technique is really helpful for the development of various species of orchids. Haploids are widely used in many agricultural and horticultural fields [10]. By using *in vitro* technique, we can develop haploid as well as diploid plants. In this paper, the production of haploid plants is being discussed.

Haploid plants are those plants that arise from gametes or gamete-like cells [11]. They do not undergo fertilization and can still generate a viable individual plant [11]. Haploid plants contain cells that have half the number of chromosomes. Haploid plants are sterile as they contain only one set of chromosomes number [12]. Haploids have only one chromosome set, that is formed after the meiosis in male or female gametes [11]. The plant is referred to as the maternal or paternal, based upon the set of chromosomes contributed by paternal or maternal side. The haploid plant is also known as monoploid because it contains only one set of chromosomes [12]. There are many advantages of haploid plants such as fixation of homozygosity, shorten breeding cycle, and high selection efficiency [13]. Haploid plants are used in the basic plant research fields such as

cytogenetics, molecular genetics, crop evolution, plant biotechnology, and traditional plant breeding [14]. Haploid plants are an effective tool for induced mutagenic and genetic transformation studies [15]. *In vitro* techniques are an efficient way to produce haploids than other methods such as interspecific hybridization or treatment with plant growth regulators, temperature, or irradiation for haploid production [16]. *In vitro* methods of haploid production include- Androgenesis which is the production of haploids by the anther or pollen culture. In androgenesis, the haploid is generated from the male gamete [6]. Androgenesis is of two types- Direct androgenesis and indirect androgenesis including parent plant selection, selection of flower bud, microspore development stage determination, surface sterilization of young flower buds, isolation of anther/microspore from buds, inoculation of anther on an appropriate medium and regeneration and recovery of the plant [6]. Gynogenesis is the production of haploids by the ovary or ovule culture [6]. In gynogenesis, the female gametes are used for the production of the haploid plant. The common steps for gynogenesis include selection of the healthy parent plant, checking of ovary/ovule developmental stage by histology, selection of appropriate flower bud and surface sterilization, inoculation of ovary/ovule in a suitable medium, incubation of ovary/ovule culture, regeneration of callus/embryo and haploid plant recovery [6].

The *in vitro* culture technique was first given by Haberlandt in 1902. *In vitro* conditions in haploid production effectively reduce the juvenile period [8,17,18]. This technique is also called tissue culture, organ culture, and cell culture [2]. Good quality and disease-free orchid plants can be obtained by using this technique [2].

The desired plant can be selected from any forest or garden. The collected plant is then replanted in a pot filled with soil [7]. This plant is kept under *in vitro* conditions. The *in vitro* conditions provide the environment that promotes the growth of the cultured plant [7]. *In vitro* conditions comprises of a proper physical and chemical environment [16].

The developmental stage of microspores is crucial. Androgenesis is the production of haploid microspore and their *in vitro* culture. This is only possible with the immature anther containing immature pollen [16]. Juvenility has important role to play in regeneration during androgenesis [19]. Pre-treatment of anthers are the most

efficient approach to the anther culture that involve a change in temperature [16]. Cold pre-treatments kill non-viable microspores and anthers [16]. It blocks the microspores in the first mitosis. Thus, change the development from gametophytic phase to sporophytic phase and these pre-treatments maintain the cellular environment which are conducive for embryogenic development [16].

Culture media is a complex solution that provides suitable conditions for cell growth. Several culture media are present. Different media are suitable for different plant species such as Murashige and Skoog (MS) medium [16], Knudson (KC) medium [7], Vacin and Went medium [7], Linsmaier and Skoog (LS) medium [20] and Nitsch and Nitsch (NN) medium [16]. Among all of these media, MS medium is the most commonly used medium. The salt composition of any medium does not affect the frequency of initial division, it affects the planting efficiency a little, but it enhances embryogenesis and regeneration process [16]. The sugar content in media also shows several effects. A high concentration of sugar is crucial for inducing androgenesis [16]. The proper growth of a plant is also affected by the growth regulators, carbon source, and growth additives. The growth regulators play an essential role in deciding the regeneration pathway to be followed by explant in nutrient regime [21].

The success of *in vitro* production depends upon the right culture conditions such as temperature and light. The ideal conditions for *in vitro* haploid production are different for different species [16]. Anthers are kept under dark conditions with a temperature of 20- 30°C [16]. The optimum temperature for the field grown material is usually 2°C which is higher than the greenhouse-grown material. *In vitro* microspore culture is all about culturing immature pollen grains that are obtained by the donor plant [22]. The methodology of microspore culture is similar to that of the anther culture [22]. This method is slower than the anther culture method for development, but the production of the embryo is higher in this method [16].

The method of *in vitro* haploid production has now been adopted in various plant reproducing programs world over. The methodology has emerged as an efficient technique for prompt harvest improvement for gene transfer, part of chromosome, and whole chromosomes [12,14,23,24]. Being economically stable

technique, it is effectively utilised in biodiversity conservation [2]. Thus, *In vitro* haploid production technique is a very productive and advisable way for haploid production [6]. Haploid plants have several applications in the fields of basic research, classical plant genetics, cytogenetics, genetic transformation research, conventional plant breeding studies, genome mapping, and many more [12].

**Significance:** *In vitro* haploid production is broadly used to produce new varieties and cultivars. The technique is also employed to conduct genetic analysis of several important agricultural crop.

### 3. CONCLUSION

Orchids are the most beautiful and commercially very valuable flowers. They are very expensive. The improvement in orchid cultivation can effectively develop the economy of our country. The natural breeding cycle of orchids is very long and time-consuming. It takes several years for orchids to blossom. *In vitro* conditions can effectively shorten the time required of breeding cycle. *In vitro* haploid production can also be used to develop viable seedlings of rare, endangered, and commercially valuable orchids. *In vitro* haploid production raises good-quality, orchid plant free of any kind of disease. Success of this technique is based upon identifying right stage of the capsule at harvesting time, the growth adjuvants, and suitable nutrient regime. *In vitro* haploid production is very useful in developing homozygous pure lines, such pure lines are utilised in various plant breeding programs, and in the research activities pertaining to plant genetics and breeding.

### CONSENT

It is not applicable.

### ETHICAL APPROVAL

It is not applicable.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Kaur Saranjeet. Biodiversity conservation with special reference to orchids. In:

- Padmavathi P, Kandru A, Das BK, Upriti BM, Mishra S, editors. Biodiversity Assessment: Tool for Conservation. 1st ed. Maharashtra: Bhumi Publishing, Kolhapur; 2017.
2. Sarmah D, Kolukunde S, Sutradhar M, Singh B, Mandal T, Mandal N. A Review on: *In vitro* Cloning of Orchids. *Int J Cur Microbiol App Sci.* 2017;6(8):1909-1927.
  3. Sipayung P, Matanari J, Lafau M, Sulastri Y, Ginting B, Sihombing D. The effect of activated charcoal dose and benzyl amino purine concentration on the growth of Orchid plantlets in Murashige and Skoog media *in vitro*. *IOP Conference Series: Earth Environ Sci.* 2017;205: 012025.
  4. Vendrame W, Faria R, Sorace M, Sahyun S. Orchid cryopreservation. *Ciencia E Agrotecnol*, 2014;38(3):213-229.
  5. Chugh S, Guha S, Rao IU. Micropropagation of Orchids: A review on the potential of different explants. *Sci Hort.* 2009;507-520.
  6. Mishra VK, Goswami R. Haploid Production in Higher Plant. *Int J Chem Biochem Sci.* 2014;1(1):2349-2724.
  7. Kaur S, Bhutani KK. *In vitro* propagation of *paphiopedilum spicerianum* (Reichb. F.) Pfitz. *Flori Orna Biotechnol.* 2013;7(1):65-70.
  8. Teixeira Da Silva JA, Kerbauy GB, Zeng S, Chen Z, Duan J. *In vitro* flowering of Orchids. *Critical Rev Biotechnol.* 2013;34(1):56-76.  
Available:<https://doi.org/10.3109/07388551.2013.807219>
  9. Hu H. *In vitro* induced haploids in wheat. In: Jain SM, Sopory SK, Veilleux RE editors, *In vitro* haploid production in higher plants. *Current Plant Science and Biotechnology in Agriculture.* Springer, Dordrecht. 1997;26.  
Available:[https://doi.org/10.1007/978-94-017-1862-2\\_3](https://doi.org/10.1007/978-94-017-1862-2_3)
  10. Pickering RA, Devaux P. Haploid production: Approaches and use in plant breeding. In: Shewry PR. ed. *Biotechnology in Agriculture No. 5. Barley: Genetics, Biochemistry, Molecular Biology and Biotechnology*, Wallingford: C.A.B. International, U.K; 1992.
  11. Britt AB, Kappu S. CenH3: An Emerging Player in Haploid Induction Technology. *Front Plant Sci.* 2016;7:357.  
DOI: 10.3389/fpls.2016.00357
  12. Basu SK, Datta M, Sharma M, Kumar A. Invited Review Article Haploid production technology in Wheat, and some selected higher plants. *Australian J Crop Sci.* 2011;5(9):1087-1093.
  13. Mishra R, Rao G. *In vitro* Androgenesis in Rice: Advantages, Constraints, and Future prospects. *Rice Sci.* 2016;23(2):57-68.
  14. Touraev A, Forster BP, Jain SM. *Advances in haploid production in higher plants.* The Netherlands: Springer Science and Business Media B.V; 2009.
  15. Folling L, Olesen A. Transformation of Wheat (*Triticum aestivum* L.), microspore-derived callus and microspores by particle bombardment. *Plant Cell Rep.* 2002;20:629-663.
  16. Atanassov A, Zagorska N, Boyadjiev P, Djilianov D. *In vitro* production of haploid plants. *World J Microbiol Biotechnol.*1995;11(4):400-408.
  17. Blanchard MM, Runkle ES. Benzylamide promotes flowering in *Doritaenopsis* and *Phalaenopsis* Orchids. *J Plant Growth Regul.* 2008;27:141-150.
  18. Taylor NJ, Staden J Van. Towards an understanding of the manipulation of *in vitro* flowering. In: Teixeira da Silva JA. ed. *Floriculture, Ornamental and Plant Biotechnology: Advances and Topical Issues.* 1st ed. IV Vol. Isleworth: Global Science Books; 2006.
  19. Datta SK, Wenzel G. Single microspore derived embryogenesis and plant formation in Barley (*Hordeum vulgare*), *Arch Zeucht.* 1998;18:125-131.
  20. Saad AIM, Elshahed AM. Plant Tissue Culture Media. In: Leva A., Rinaldi LMR, eds. *Recent advances in plant in vitro culture.* Winchester: In Tech; 2012.
  21. Hussain A, Qarshi IA, Nazir H, Ullah I. Plant tissue culture: Current status and opportunities. In: Leva A, ed. *Recent advances in plant in vitro culture.* Italy: Intech Open; 2012.  
DOI: 10.5772/50568
  22. Rajcan I, Boersma JG, Shaw EJ. *Plant Genetics Techniques: Plant Breeder's Toolbox.* *Comprehensive Biotechnology.* 2nd ed.4<sup>th</sup>; 2011.  
Available:<https://doi.org/10.1016/B978-008-088504-9.00252-X>
  23. Baenziger PS, DePauw RM. Wheat breeding: Procedures and strategies. In: Carver BF. ed. *Wheat: Science and Trade.* IA, USA: Wiley-Blackwell Publishing; 2009.

24. Ceoloni C, Jauhar PP. Chromosome engineering of the Durum Wheat genome: Strategies and applications of potential breeding value. In: Singh RJ, Jauhar PP eds. Genetic resources, chromosome engineering, and crop improvement. Boca Raton, FL: CRC Press/Taylor and Francis. 2006;2.

---

© 2021 Jaswal and Kaur; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/77563>