

## **Efficacy of Herbal Medicine Administration using Neem (*Azadirachta indica* L) Leaves as Case Study**

**Duduyemi Oladejo <sup>a\*</sup>, Patinvoh Regina <sup>a</sup>, Adelowo Aisha <sup>a</sup> and J. G. Akinbomi <sup>a</sup>**

<sup>a</sup> Department of Chemical Engineering, Lagos State University, Epe, Lagos State, Nigeria.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/CSJI/2021/v30i1130262

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/78460>

**Original Research Article**

**Received 06 October 2021  
Accepted 12 December 2021  
Published 15 December 2021**

### **ABSTRACT**

Plants used for treatment of diverse ailments primitively are concocted and used indiscriminately. The efficacy of "Herbal medicine" which is an ancient tradition, used in some parts of Nigeria was investigated to establish herbal applications of Neem (*Azadirachta indica*) leaves. Neem leaves (*Azadirachta indica*) was popularly recognised for the treatment of malaria fever. The leaves were plucked at appropriate peak period of the day when oil and moisture contents were recorded maximum. The peak period of contents was found to occurred between 8am-10am on a very sunny days, 10am to 12 on cool days, and 3:00pm to 5:00pm on cloudy days. Aqueous extract from macerated Neem leaves was subjected to qualitative and quantitative analysis. Available phytochemicals evaluated include; saponin (34.89mg/g), tannin (31.715mg/g), flavonoid (31.835mg/g), phenol (43.59mg/g), terpernoid (14.585mg/g), cardiac glycosides (39.335mg/g), steroid (16.185mg/g) and alkaloid (28.76mg/g). These values differ significantly to recommended oral dosage formulation for human consumption: Saponin (1.433ml), Tannin (1.418ml), Flavonoid (13.91ml), Phenol (2.29ml), Terpernoid (8.23ml), Cardiac Glycosides (0.003177ml), Steroid (0.62mg/g). Consequently, local consumption of herbal resources should be regulated to avoid abuse and long or short-term effects of drug contents as proven in the neem leaves as local herbs.

**Keywords:** *Phytochemicals; calibration; dosage; peak periods; herbal; concocted.*

\*Corresponding author: E-mail: [Engrduduyemi@yahoo.com](mailto:Engrduduyemi@yahoo.com), [oladejo.duduyemi@lasu.edu.ng](mailto:oladejo.duduyemi@lasu.edu.ng);

## 1. INTRODUCTION

The use of herbs for medicinal purpose is indigenous but the indiscriminate usage of herbal products by the people had become habitual in some localities because of their believes. This had become worrisome in recent years as people engage in haulage and hawking of diverse mixtures of herbal leaves, barks of trees and roots as well as some fruits without due consideration of the dosage and the long-term effects of the contents and its accumulation in the body system. Medicinal plants are used by humans to treat sickness [1]. Demand for medicinal herbs is increasing quickly, owing mostly to the negative side effects of synthetic chemical medicines, believed ineffectiveness of drugs in some cases, poverty, lack of purchasing power, ignorance and addiction to available options. The global demand for natural ingredients encourages the cultivation of therapeutic crops on a local scale while regulated and sustainable collection of wild plants have not been given serious consideration. Such efforts have the potential to expand rural employment in developing nations, promote global commerce, and improve the lives of millions of people [2]. The chemical compounds from plant parts create healing effects in human body, the most important of these plant bioactive components are alkaloids, tannins, flavonoids and phenolic compounds [3].

Plants are good source of available and low-cost medicine. Medicinal plants are therapeutically beneficial, yet they have little or no adverse effects [4]. Plants parts and their morphologies such as roots, stems, barks, leaves, berries, fruits, seeds, and exudates, comprise essentially significant components employed in the medication due to their low side effects and low cost. Medicinal plant extracts and products as natural resources have divers characteristics and so are widely acknowledged to be the source of many pharmacologically active medications including analgesics, antidiuretics, stimulant and antibiotics. Nigeria is endowed with a vast array of these plant species, this heritage provides for her people food, medicine, and other necessities [4].

Some of the available medicinal plant parts engaged for use include; Pawpaw leaves, Mango leaves, Lemongrass, Chamomile, Eucalyptus, Aloe Vera, Scent Leaf, Butterleaf, Neem, Gingko, Teatree, Lavender, Moringa, Candle Plant, Wild Lettuce, Mint leaves, Water leave, Avocado

leaves [5]. These plants are administered in divers' recipes either as single or at times in selected combinations for different ailment.

*Azadirachta indica* (Neem) is a tree that is evergreen. It is a member of the Meliaceae family, it plays an important role in health-promoting effect because it is a rich source of antioxidant. It has a complex of distinct composition such as Nimbin, Nimbidine, Nimbolide and Limonoids and these ingredients play a part in the treatment of diseases. Quercetin and  $\beta$ -sitosterol became the first polyphenols Flavonoids isolated from fresh neem leaves and considered to have antifungal and antibacterial function [6]. Neem has various biological and pharmacological effects such as antibacterial, antifungal, anti-inflammatory, antibiotics, antigastric ulcer antitumour activities, antiarthritic, antipyretic, hypoglycemic, antigastric ulcer, antimalarial [7]. However, the qualitative analysis and application of this findings has not been put into use as local applications evolves around modulation of multiple constituents of different parts of medicinal plants concocted and used indiscriminately. It is enthused with a mindset that it can cure all forms of sickness as 'Gbogbonise' in some local settings in the South-West zone of Nigeria. Therefore, the objective of this work is to investigate the usage of neem leave extract in relevance to recommended dosage for human consumption.

## 2. MATERIALS AND METHODS

### 2.1 Preliminary Investigations

Questionnaires were prepared to investigate the most widely sourced medicinal leaves used for treatment of most common ailment in the local areas of Epe town suburb of Lagos State Nigeria. Investigation suggested (Malaria) and locally sourced alternative medicine (Herbal) includes the leaves were Mango, Pawpaw and Neem leaves. The most recommended herb by the data analysis using five hedonic scale for the purpose of this research is Neen. The appropriate time for cutting or harvesting the leaves was investigated using Oil-Moisture Analyser to evaluate the oil-water content of the leaves while on the stalk. The leaves contents were recorded hourly to obtain periods of maximum oil-water percent (peaks) between the hours of 8:00 am and 5:00 pm daily while Average recorded data were determined.

## 2.2 Sample Preparations

Fresh leaves free of contaminant and perforations were collected during the peak hours of the day between 10am-12pm when oil and moisture contents were at high percentage using an oil and moisture content analyser. The freshly collected neem leaves was dried at room temperature  $25\pm 2^{\circ}\text{C}$  until it was bone dried and stored in a desiccator in preparation for extraction (Biswas .et al, 2002). About 500g of the dried leaves was weighed and macerated to break cell wall to release soluble phytochemicals when dissolved. The macerated sample was soaked in 2000 ml (98% Laboratory grade) ethanol to maintain a liquor ratio of 4:1 volume to weight [8]. The media is frequently agitated every 3 hours at room temperature for a period of three days until the soluble matter properly dissolved. Lowering particle size increases surface contact between samples and extraction solvents. The extract from the macerated neem was collected by filtration using Whatman filter paper and subjected to further analysis. A sample of the fresh Neem leaves and the bone dried samples are shown in plate 1.

### 2.2.1 Analysis of phytochemical contents

Qualitative analysis was carried out to ascertain the presence of the different phytochemical compounds contained in the extract. The extracts were subjected to thin-layer and paper chromatographic procedures at the laboratory. This was done in order to separate the components into individual compounds for appropriate identification of all components of the extracts. Preliminary phytochemical analysis of the extracts was carried out with reference to the standard methods [9].

### 2.3 Estimation of Alkaloids

Total alkaloid was measured using methods as described by [10] with slight modification. 1ml plant extract, 5 ml pH 4.7 phosphate Buffer was added and 5 ml BCG solution and the mixture were shake with 4 ml of chloroform. The extracts were collected in a 10-ml volumetric flask and then diluted to adjust volume with chloroform. The absorbance of the complex in chloroform was measured at 470 nm against blank prepared as above but without extract. Atropine is used as a standard (using calibration curve) and compared the assay with Atropine equivalents.

Preparation of Standard Curve Accurately measure aliquots (0.4, 0.6, 0.8, 1 and 1.2 mL) of atropine standard solution and transfer each to

different separatory funnels. Add 5 mL pH 4.7 phosphate buffer and 5 mL BCG solution. Shake mixture with 1, 2, 3 and 4 mL of chloroform. The extracts were collected in a 10 mL volumetric flask and then diluted to volume with chloroform. The absorbance of the complex in chloroform was measured at 470 nm against blank prepared as above but without atropine.

### 2.4 Estimation of Steroids

Total steroid was measured using methods as described by [11] with slight modification. 1ml of test extract of steroid solution was transferred into 10ml volumetric flasks, Sulphuric acid (4N, 2ml) and iron (III) chloride (0.5% w/v, 2 ml), were added, followed by potassium hexacyanoferrate (III) solution (0.5% w/v, 0.5 ml). The mixture was heated in a water-bath maintained at  $70\pm 2^{\circ}\text{C}$  for 30 minutes with occasional shaking and diluted to the mark with distilled water. The absorbance was measured at 780 nm against the reagent blank. Total steroids in extracts were expressed in terms of cholesterol equivalents (mg of CHO/g of extract).

### 2.5 Estimation of Flavonoids

Aluminum chloride colorimetric method (Chang *et al.*,2002) with some modifications was used to determine flavonoid content. Plant extract (1mL) in methanol was mixed with 1ml of methanol, 0.5 mL aluminum chloride (1.2%) and 0.5 mL potassium acetate (120 mM). The mixture was allowed to stand for 30 min at room temperature; then the absorbance was measured at 415 nm. Quercetin was used as standard. Flavonoid content is expressed in terms of quercetin equivalent (mg g<sup>-1</sup> of extracted compound).

### 2.6 Estimation of Total Phenols

The Follins method described by (Pearson 1976) was used to determine the phenol content. 1ml of the extract was placed in a test tube, 2.5ml of 10% Folin-Ciocalteu's reagent dissolved in water and 2.5 ml of 7.5% of NaHCO<sub>3</sub> aqueous solution were added. The samples were thereafter incubated in a thermostat at 45°C for 45 min. The absorbance was determined using spectrophotometer at wave length = 765 nm. The samples were prepared in triplicate for each analysis and the mean value of absorbance was obtained. The same procedure was repeated for the standard solution of gallic acid and the calibration line was construed. Based on the measured absorbance, the concentration of gallic acid equivalent expressed in terms of (mg of GA/g of extract).



**Plate 1a. freshly collected neem leave**



**Plate 1b. Bone dried Neem leaves samples**

## 2.7 Estimation of Tannins

Tannin content was determined by the Folin-Denis colorimetric method described by Kirk and Sawyer [12]. 5g sample was dispersed in 50mls of distilled water and shaken. The mixture was allowed to stand for 30min at 28°C before it was filtered through Whatman No. 42 grade of filter paper. 2ml of the extract was dispersed into a 50ml volumetric flask. Similarly, 2ml standard tannin solution (tannic acid) and 2ml of distilled water were put in separate volumetric flasks to serve as standard and reagent was added to each of the flask and the 2.5ml of saturated  $\text{Na}_2\text{CO}_3$  solution added. The content of each flask was made up to 50ml with distilled water and allowed to incubate at 28°C for 90 min. Their respective absorbance was measured in a spectrophotometer at 765nm using the reagent blank to calibrate the instrument at zero.

## 2.8 DNS reducing sugar assay

This assay is based on the protocol described by Hussain *et al.*, [13] with slight modification; The 3,5-Dinitrosalicylic acid (DNS) reagent was prepared by mixing 1.6 g NaOH and 1.0 g dinitro salicylic acid (Sigma) in 70 mL dH<sub>2</sub>O and the mixture was heated in boiling water to dissolve. Once dissolved, 3.0 g  $\text{Na}_2\text{K}$  tartrate (Sigma) was added to the solution and swirled until dissolved followed by further addition of H<sub>2</sub>O to make up to 100 mL. The reagent was stored dark at room temperature. 1ml of the plant samples is taken and 3ml of DNSA was added and boiled for 10min and absorbance was taken at 540nm in an

Atomic Absorption Spectrum equipment. Glucose was used as standard (100mg/ml).

## 2.9 *In vitro* Antioxidant Assays reducing Power Assay [14]

The reducing power was determined according to the method with slight modifications. Reaction was carried out in a mixture containing 1 ml of sample (25-100  $\mu\text{g}/\text{ml}$ ), 2.5 ml of 0.1 M sodium phosphate buffer (pH 6.6) and 2.5 ml of 1%, w/v potassium ferro cyanate [ $\text{K}_3\text{Fe}(\text{CN})_6$ ] by incubating at 50°C for 20 min. After addition of 2.5 ml trichloroacetic acid (10%, w/v), the mixture was centrifuged at 5000rpm for 10 min. The upper layer (5 ml) was mixed with 0.5 ml of fresh  $\text{FeCl}_3$  (0.1%, w/v), and the absorbance at 700 nm was measured against a blank. Gallic acid was used as the control.

## 2.10 DPPH radical scavenging assay [15]

Briefly 0.1 mM solution of DPPH in ethanol was prepared; 1ml of the solution was added to 1 ml of extract in water at different concentrations (25-100  $\mu\text{g}/\text{ml}$ ). The mixture was vigorously shaken and afterwards allowed to stand at room temperature for 30 min to settle. Then the absorbance was measured at 517 nm by using a UV-Visible Spectrophotometer. When of the reaction mixture was subjected to Lower absorbance, it indicated higher free radical scavenging activity. The percent DPPH scavenging effect was calculated using equation 1;

$$\text{DPPH Scavenging effect (\%)} = \frac{A_0(\text{Control}) - A_1(\text{Sample})}{A_0(\text{Control})} \times 100 \quad (1)$$

Where  $A_0$  was the absorbance of the control and  $A_1$  was the absorbance in the presence of the standard sample or extract. The  $IC_{50}$  value represented the concentration of the compounds that caused 50% inhibition of DPPH radical formation.

### 2.10.1 Hydroxyl radical scavenging assay [16]

The mixture containing 1 ml of sample (0.1–0.5 mg/ml), 1 ml of 9 mM  $FeSO_4$  and 1 ml of 0.3%  $H_2O_2$  in 0.5 ml of 9 mM salicylic acid–ethanol solutions was shaken vigorously and incubated at 37°C for 30 min. Then, the absorbance of the reaction mixture was determined at 510 nm. Gallic Acid (GA) was used as the positive control. The hydroxyl radical scavenging activity was calculated by equation 2;

$$\text{Scavenging Activity (\%)} = \left[ 1 - \frac{A_1 - A_2}{A_0} \times 100 \right] \quad (2)$$

Where  $A_0$  is the absorbance of the control (ethanol instead of sample);  $A_1$  is the absorbance of the sample, and  $A_2$  is the absorbance of the sample only (salicylic acid–ethanol solution instead of  $FeSO_4$  and  $H_2O_2$  solutions). The  $IC_{50}$  value represented the concentration of the compounds that caused 50% inhibition of hydroxyl radical formation.

### 2.10.2 Superoxide radical scavenging assay [16]

Reaction was carried out in a mixture containing 4.5 ml of 50 mM Tris–HCl buffer (pH 8.2), 0.4 ml of 25 mM pyrogallol solution and 1 ml of sample (0.5–5 mg/mL) by incubating at 25°C for 5 min. Finally, 1 ml of 8 mM HCl solution was dripped into the mixture promptly to stop further reaction. The absorbance of the mixture was measured at 420 nm frequency. Gallic Acid was used as the positive control. The superoxide radical scavenging activity was valued using equation 3;

$$\text{Scavenging activity (\%)} = \left[ 1 - \frac{A_1 - A_2}{A_1} \times 100 \right] \quad (3)$$

Where  $A_0$  is the absorbance of the control (water instead of sample),  $A_1$  is the absorbance of the sample, and  $A_2$  is the absorbance of the sample only (Tris–HCl buffer instead of pyrogallol solution). The  $IC_{50}$  value represented the concentration of the compounds that caused 50% inhibition of superoxide radical formation.

### 2.10.3 Hydrogen peroxide radical scavenging assay [1]

Reaction mixture containing 1 ml of sample (0.1–2 mg/mL), 2.4 ml of phosphate buffer (0.1 M, pH 7.4) and 0.6 ml of  $H_2O_2$  solution (40 mM) was shaken vigorously and incubated at room temperature for 10 min. Then, the absorbance of the reaction mixture was determined at 230nm. Gallic Acid was used as the positive control. The  $H_2O_2$  scavenging activity was calculated as follows:

Where  $A_0$  is the absorbance of the control (water instead of sample),  $A_1$  is the absorbance of the sample, and  $A_2$  is the absorbance of the sample only (phosphate buffer instead of  $H_2O_2$  solution). The  $IC_{50}$  value represented the concentration of the compounds that caused 50% inhibition of  $H_2O_2$ . It is calculated by equation 4;

$$\text{Scavenging activity (\%)} = \left[ 1 - \frac{A_1 - A_2}{A_0} \times 100 \right] \quad (4)$$

### 2.10.4 $\beta$ -carotene linoleate bleaching assay [17]

In the assay, linoleic acid produces hydroperoxides (ROS), and this oxidizes the  $\beta$ -carotene at 50°C. The presence of antioxidants in the extract will minimize the oxidation of  $\beta$ -carotene by hydroperoxides. Hydroperoxides formed in this system will be inactivated by antioxidants from extracts. Gallic acid was used as the standard. B- Carotene (0.2 mg in 1 ml of chloroform), linoleic acid (0.02 ml), and Tween 80 (0.2 ml) were transferred into a round-bottomed flask. Chloroform was removed at room temperature using a rotary evaporator. Following evaporation, 50 ml of distilled water was added to the mixture and shaken vigorously to form an emulsion. Two milliliters of aliquots of the emulsion was pipetted into test tubes containing of mushroom extracts (different concentration of 20–100  $\mu$ g/ml) and immediately placed in a water bath at 50°C. The absorbance was read at 20 min intervals for 2 h at 470 nm using UV-vis spectrophotometer. The antioxidant activity (I) was expressed as a percent of inhibition relative to the control, using equation 5 below;

$$\text{Inhibition (\%)} = \frac{A_c - A_s}{A_c} \times 100 \quad (5)$$

Where  $A_c$  and  $A_s$  represent the bleaching rates of  $\beta$ -carotene without and with the addition of antioxidant, respectively.

### 2.10.5 Lipid peroxidation Assay [18]

The reaction mixture contained 1 mL of fowl egg yolk emulsified with phosphate buffer (pH 7.4) to obtain a final concentration of 25 g/l, sample (different concentration of 20–100 µg/ml), and 100 µL of 1000 µM FeCl<sub>2</sub>. The mixture was incubated at 37°C for 1 h before being treated with 0.5 ml of freshly prepared 15% trichloroacetic acid (TCA) and 1.0 ml of 1% thiobarbituric acid (TBA). The reaction tubes were further incubated in boiling water bath for 10 min. Once cooled to room temperature, the tubes were centrifuged at 3500 rpm for 10 min to remove precipitated protein. The absorbance at 532 nm was determined spectrophotometrically. Gallic acid was used as positive control. The percentage inhibition was calculated from equation 6;

$$\text{Inhibition (\%)} = \frac{A_b - A_s}{A_b} \times 100 \quad (6)$$

Where  $A_b$  is the absorbance of the blank without the extract or ascorbic acid and  $A_s$  is the absorbance in the presence of the extract or gallic acid

### 2.10.6 Inhibition percentage (Griess Reagent)

Sodium nitroprusside (10mM) in phosphate buffer saline was mixed with different concentrations (25-100 µg/ml) of each extract, and incubated for 30min, after incubation period, 0.5ml of Griess reagent (1% sulfanilamide, 2% H<sub>2</sub>PO<sub>4</sub> and 0.1% N-(1-naphthyl) ethylenediamine dihydrochloride) was added. Absorbance was taken at 550nm and ascorbic acid was used as standard. IC<sub>50</sub> which is an inhibitory concentration of each extract required to reduce 50% of nitric oxide formation was determined. The same reaction mixture without the extract but equivalent amount of methylated spirit was used as control. It is calculated using equation 7;

$$\text{Nitric oxide (inhibition \%)} = \frac{A(\text{control}) - A(\text{sample})}{A(\text{control})} \times 100 \quad (7)$$

## 3. RESULTS AND DISCUSSION

### 3.1 Preliminary Investigations

Questionnaires distributed to local merchants and users of herbs was analysed to investigate three medicinal leaves commonly taken for

treatment of ailment such as malaria in EPE, suburban areas of Lagos State. The neem leaf was adjudged widely used for the treatment of malaria sickness either singly or along other recipes. The investigation showing the distribution of opinion poll on the uses of neem leaves choice and applications as shown in Fig. 1.

### 3.2 Oil and Moisture Content Analysis of Sampled leaves

The oil-moisture detector is used to analyse content of the leaves while still on the stalk hourly for five days between 8:00 am - 5:00 pm to determine the best time for sample collection. This occurred when the oil and water content recorded are practically at the peak. It was observed that the trend of oil and water movement along the plant morphology followed the same pattern of variation. highest and almost at the same level. In this regard, the right time for the harvest of leaves were found to be between 8am and 10am on early sun rise day, and on a fair weather day between 10 am and 12:00 Noon but on a cool days perhaps with rain, ideal time was between 3pm to 5pm as shown in Fig. 2 Fig. 3 and Fig. 4 respectively.

The distributions of oil and moisture content in the neem leaves depends on the Evapo-transpiration phenomenon associated with the heating effects of the sun rays and the osmotic pulling of the nutrients between the morphology of the plants transport system. This is evidenced by the peaks recorded in the repeated days with similar weather description. Therefore, the time of the day and the prevailing weather condition is very important in obtaining contents and quantity of herbal resources.

### 3.3 Analysis of Extract

The group phytochemicals detected in the neem leaves as well as the amounts of each based on the qualitative and quantitative analysis of extract on the basis 100g macerated neem leaves is presented in Table 1.

### 3.4 Required Dosage Recommendation

The recommended dosage of phytochemicals for human consumption was evaluated according to Toney-Butler et al (2021) as shown in equation 8:

$$\text{Dosage} = \frac{D \times Q}{H} \quad (8)$$




Fig 1. Analysis of questionnaire distribution




Fig 2. Plot of oil and moisture content for day1 (Early Sunrise Day)




Fig. 3. Plot of and moisture content for day 2 (Normal bright day)




**Table 1. Quantitative analysis of Neem contents**

	<b>Saponin mg/100g</b>	<b>Tannin mg/100g</b>	<b>Flavonoid mg/100g</b>	<b>Phenol mg/100g</b>	<b>Alkaloids mg/100g</b>	<b>Reducing sugar mg/100g</b>	<b>Terpernoid mg/100g</b>	<b>Cardiac glycoside mg/100g</b>	<b>Steroid mg/100g</b>
Sample1	34.56	31.57	31.97	43.40	29.38	33.93	14.30	39.54	15.73
Sample2	35.22	31.86	31.70	43.79	28.14	32.99	14.87	39.13	16.64
Average	34.89	31.715	31.835	43.59	28.76	33.46	14.585	39.335	16.185

**Table 2. Required oral dosage**

<b>Phytochemicals</b>	<b>Average quantity mg/g</b>	<b>D (prescribed intake) mg/g</b>	<b>Dosage (ml)</b>
Saponin	34.89	50mg/g/day(Diwan., et al 2000)	1.433
Tannin	31.715	45mg/g/day[19]	1.418
Flavonoid	31.835	443mg/g/day(Tresserra., et al 2013)	13.91
Phenol	43.59	100mg/g/day[20]	2.29
Terpernoid	14.585	120mg/g/day[21]	8.23
Cardiac glycosides	39.335	0.125mg/d/day[22]	0.003177
steroid	16.185	10mg/g/day[23]	0.62



**Fig. 4. Plot of oil and moisture content for day 3 (Raining day)**



Where; D – prescribed intake

H – available amount

Q – quantity(1ml)

The available quantities and prescribed dosages for different phytochemicals and flavonoids are presented in Table 2. It revealed the disparity between available content and for references and comparison with the dosage indiscriminately consumed by herbal product users especially of the neem origin the respective.

#### 4. CONCLUSION AND RECOMMENDATIONS

The oil and moisture content in which the phytochemicals, active ingredients and transport systems are contained was seen to follow the same pattern though in different quantity, availability of phytochemicals in oil and moisture is weather dependent, oral dosage required for human consumption has been compared to the orthodox prescription. The disparity calls for cautions as daily consumption of herbal medicine is increasing as herbal and advocates of herbal medicines and practitioner are increasing. The correct utilization of these products in the right dosages should therefore be regulated as overdose and incorrect or incompatible combinations might have serious consequences on the short or later in the future of consumers.

Medicinal plants in general, and neem in particular, are a gift from nature, and their usages should be encouraged, particularly in the production of pharmaceuticals, which provides an incentive to alternative medicine and practices. Several contemporary medications have been discovered to have traditional origins; hence, medicinal plants have been recognized. Alternative medicine therefore should be a symbiotic relationship between traditional medical practice and its orthodox counterpart in preventing health complications. Dosage of local herbs should be controlled and contents regulated, to avoid abuse and long-time consequential effects of accumulated phytochemicals.

#### NOTE

The study highlights the efficacy of "Herbal medicine" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Sarbani Dey Ray. Reducing power and superoxide radical scavenging activity of a triterpene derivative isolated from root bark of *Zizyphus nummularia* Aubrev. *Asian Journal of Pharmacy and Pharmacology*;2019. Available:<https://doi.org/10.31024/ajpp.2019.5.1.21>.
2. Anita M. Medicinal and aromatic plants: Monitoring the effectiveness of biological conservation;2004.
3. Hill RA. Terpenoids. In Thomson RH, (ed). *Chemistry of Natural Products*, Blackie Academic and Professional. London. 1985;106-134.
4. Odugbemi T, Akinsulire O. Medicinal plants by species names 'in: Odugbemi T, editor: 'Outlines and pictures of medicinal plants from Nigeria', University of Lagos Press. 2006;95-156.
5. Monier M. Abd El-Ghani. *Traditional medicinal plants of Nigeria*;2016.
6. Govindachari TR, Suresh G, Gopalakrishnan G, Banumathy B, Masilamani S. Identification of antifungal compounds from the seed oil of *Azadirachta indica*. *Phytoparasitica*. 1998;26(2):109–116. DOI: 10.1007/bf02980677. [CrossRef]
7. Bandyopadhyay U, Biswas K, Sengupta A, et al. Clinical studies on the effect of Neem (*Azadirachta indica*) bark extract on gastric secretion and gastroduodenal ulcer. *Life Sciences*. 2004;75(24):2867–2878. DOI: 10.1016/j.lfs.2004.04.050. [PubMed] [CrossRef]
8. Handa SS, Khanuja SPS, Longo G, Rakesh DD. *Extraction Technologies for Medicinal and Aromatic Plants*, (1stedn), no. 66. Italy: United Nations Industrial Development Organization and the International Centre for Science and High Technology;2008.
9. Sofowora A. *Medicinal Plants and Traditional Medicine in Africa*. Spectrum Books Ltd., Ibadan, Nigeria. 1993;191-289.
10. Fazel Shamsa, Hamidreza Monsef, Rouhollah Ghamooshi and Mohammadreza Verdianrizi. Spectrophotometric determination of total

- alkaloids in some Iranian medicinal plants;2008.
11. Salomi P, Sucharitha B, Swathi T, Hemalatha P, Ravivarma K, Venkata Ramana A, Ravindra Reddy K. Colorimetric method for determination of corticosteroids by UV Visible Spectroscopy and its application to Ayurvedic formulations;2019.
  12. Kirk HR. Sawyer Fraith Pearson Chemical Analysis of Food (8th edition), Longman Scientific and Technical, Edinburgh. 1998;211-212
  13. Hussain T, Shah SZA, Zhao D, Sreevatsan S, Zhou X. The role of IL-10 in *Mycobacterium avium* ssp. *paratuberculosis* infection Cell Commun. Signal. 2016;14:2927905994
  14. Oyaizu M. Studies on products of browning reaction prepared from glucosamine, Jpn J of Nutr. 1986;44:307–15.
  15. Shimada K, Fujikawa K, Yahara K, Nakamura T. Antioxidative properties of xanthone on the auto oxidation of soybean in cyclodextrin emulsion. J. Agr. Food Chem. 1992;40:945–948.
  16. Zhang SB, Wang Z, Xu SY. Antioxidant and antithrombotic activities of rapeseed peptides. Journal of the American Oil Chemists’ Society, 2008;85:521–527.
  17. Cheung LM, Cheung PCK, Ooi VEC. Antioxidant activity and total phenolics of edible mushroom extracts. Food Chem. 2003;81:249-55.
  18. SinanInce AN, Ismail K, Ibrahim HC, Fatih FA, Abdullah E. The effects of dietary boric acid and borax supplementation on lipid peroxidation, antioxidant activity, and DNA damage in rats. Journal of Trace Elements in Medicine and Biology. 2010;24:161–164.
  19. Samanta S, Giri S, Parua S, Nandi DK, Pati BR, Mondal KC. Impact of tannic acid on the gastrointestinal microflora, Microbial Ecology in Health and Disease, 2004;16(1):32-34.  
DOI: 10.1080/08910600310026158
  20. WHO. IPCS Health and Safety Guide No. 88? Published by WHO. Printed by Wissenschaftliche Verlagsgesellschaft, Stuttgart;1994.
  21. Jessica A Miller, Julie E Lang, Michele Ley, RayNagle, Chiu-Hsieh Hsu, Patricia A Thompson, Catherine Cordova, Amy Waer, H-H Sherry Chow. Human breast tissue disposition and bioactivity of limonene in women with early stage breast cancer;2013.
  22. John P. Cunha, DO, FACOEP 2020 Digoxin.
  23. Conn DL Resolved: low-dose prednisone is indicated as a standard treatment in patients with rheumatoid arthritis.

© 2021 Oladejo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/78460>